

Descent distribution on Catalan words avoiding a pattern of length at most three

Jean-Luc Baril, Sergey Kirgizov and Vincent Vajnovszki
LE2I, Université de Bourgogne Franche-Comté
B.P. 47 870, 21078 DIJON-Cedex France
e-mail: {barjl,sergey.kirgizov,vvajnov}@u-bourgogne.fr

March 18, 2018

Abstract

Catalan words are particular growth-restricted words over the set of non-negative integers, and they represent still another combinatorial class counted by the Catalan numbers. We study the distribution of descents on the sets of Catalan words avoiding a pattern of length at most three: for each such a pattern p we provide a bivariate generating function where the coefficient of $x^n y^k$ in its series expansion is the number of length n Catalan words with k descents and avoiding p . As a byproduct, we enumerate the set of Catalan words avoiding p , and we provide the popularity of descents on this set. Some of the obtained enumerating sequences are not yet recorded in the On-line Encyclopedia of Integer Sequences.

Keywords: Enumeration, Catalan word, pattern avoidance, descent, popularity.

1 Introduction and notation

Combinatorial objects counted by the Catalan numbers are very classical in combinatorics, with a variety of applications in, among others, Biology, Chemistry, and Physics. A length n *Catalan word* is a word $w_1 w_2 \dots w_n$ over the set of non-negative integers with $w_1 = 0$, and

$$0 \leq w_i \leq w_{i-1} + 1,$$

for $i = 2, 3, \dots, n$. We denote by \mathcal{C}_n the set of length n Catalan words, and $\mathcal{C} = \cup_{n \geq 0} \mathcal{C}_n$. For example, $\mathcal{C}_2 = \{00, 01\}$ and $\mathcal{C}_3 = \{000, 001, 010, 011, 012\}$. It is well known that the cardinality of \mathcal{C}_n is given by the n th Catalan number $\frac{1}{n+1} \binom{2n}{n}$, see for instance [12, exercise 6.19.u, p. 222], which is the general term of the sequence [A000108](#) in the On-line Encyclopedia of Integer Sequences (OEIS) [11]. See also [9] where Catalan words are considered in the context of the exhaustive generation of Gray codes for growth-restricted words.

A *pattern* p is a word satisfying the property that if x appears in p , then all integers in the interval $[0, x - 1]$ also appear in p . We say that a word $w_1 w_2 \dots w_n$ contains the pattern $p = p_1 \dots p_k$ if there is a subsequence $w_{i_1} w_{i_2} \dots w_{i_k}$ of w , $i_1 < i_2 < \dots < i_k$, which is order-isomorphic to p . For example, the Catalan word 01012312301 contains seven occurrences of the pattern 110 and four occurrences of the pattern 210. A word *avoids* the pattern p whenever it does not contain any occurrence of p . We denote by $\mathcal{C}_n(p)$ the set of length n Catalan words avoiding the pattern p , and $\mathcal{C}(p) = \cup_{n \geq 0} \mathcal{C}_n(p)$. For instance, $\mathcal{C}_4(012) = \{0000, 0001, 0010, 0011, 0100, 0101, 0110, 0111\}$, and $\mathcal{C}_4(101) = \{0000, 0001, 0010, 0011, 0012, 0100, 0110, 0111, 0112, 0120, 0121, 0122, 0123\}$. For a set of words, the *popularity* of a pattern p is the overall number of occurrences of p within all words of the set, see [4] where this notion was introduced, and [1, 7, 10, 2] for some related results.

A *descent* in a word $w = w_1 w_2 \dots w_n$ is an occurrence $w_i w_{i+1}$ such that $w_i > w_{i+1}$. Alternatively, a descent is an occurrence of the *consecutive* pattern 10 (*i.e.*, the entries corresponding to an occurrence of 10 are required to be adjacent). We denote by $d(w)$ the number of descents of w , thus the popularity of descents on a set S of words is $\sum_{w \in S} d(w)$. The distribution of the number of descents has been widely studied on several classes of combinatorial objects such as permutations and words, since descents have some particular interpretations in fields as Coxeter groups or theory of lattice paths [3, 6].

The main goal of this paper is to study the descent distribution on Catalan words (see Table 1 for some numerical values). More specifically, for each pattern p of length at most three, we give the distribution of descents on the sets $\mathcal{C}_n(p)$ of length n Catalan words avoiding p . We denote by $C_p(x, y) = \sum_{n, k \geq 0} c_{n, k} x^n y^k$ the bivariate generating function for the cardinality of words in $\mathcal{C}_n(p)$ with k descents. Plugging $y = 1$

- into $C_p(x, y)$, we deduce the generating function $C_p(x)$ for the set $\mathcal{C}_n(p)$, and
- into $\frac{\partial C_p(x, y)}{\partial y}$, we deduce the generating function for the popularity of

descents in $\mathcal{C}_n(p)$.

From the definition at the beginning of this section it follows that a Catalan word is either the empty word, or it can uniquely be written as $0(w' + 1)w''$, where w' and w'' are in turn Catalan words, and $w' + 1$ is obtained from w' by adding one to each of its entries. We call this recursive decomposition *first return decomposition* of a Catalan word, and it will be crucial in our further study. It follows that $C(x)$, the generating function for the cardinality of \mathcal{C}_n , satisfies:

$$C(x) = 1 + x \cdot C^2(x),$$

which corresponds precisely to the sequence of Catalan numbers.

We conclude this section by explaining how Catalan words are naturally related to two classical combinatorial classes counted by the Catalan numbers.

Catalan words vs. Dyck words

A *Dyck word* is a word over $\{u, d\}$ with the same number of u 's and d 's, and with the property that all of its prefixes contain no more d 's than u 's. Alternatively, a Dyck word can be represented as a lattice path starting at $(0, 0)$, ending at $(2n, 0)$, and never going below the x -axis, consisting of up steps $u = (1, 1)$ and down steps $d = (1, -1)$. There is a direct bijection $\delta \mapsto w$ between the set of Dyck words of semilength n and \mathcal{C}_n : the Catalan word w is the sequence of the lowest ordinate of the up steps u in the Dyck word δ , in lattice path representation. For instance, the image through this bijection of the Dyck word $udvudvuddvuddd$ of semilength 7 is $0011212 \in \mathcal{C}_7$. Note that the above bijection gives a one-to-one correspondence between occurrences of the consecutive pattern ddu in Dyck words and descents in Catalan words.

Catalan words vs. binary trees

In [8] the author introduced an integer sequence representation for binary trees, called *left-distance sequence*. For a binary tree T , let consider the following labeling of its nodes: the root is labeled by 0, a left child by the label of its parent, and a right child by the label of its parent, plus one. The left-distance sequence of T is obtained by covering T in inorder (*i.e.*, visit recursively the left subtree, the root and then the right subtree of T) and collecting the labels of the nodes. In [8] it is showed that, for a given length,

the set of left-distance sequences is precisely that of same length Catalan words. Moreover, the induced bijection between Catalan words and binary trees gives a one-to-one correspondence between descents in Catalan words and particular nodes (left-child nodes having a right child) in binary trees.

The remainder of the paper is organized as follows. In Section 2, we study the distribution of descents on the set \mathcal{C} of Catalan words. As a byproduct, we deduce the popularity of descents in \mathcal{C} . We consider also similar results for the obvious cases of Catalan words avoiding a pattern of length two. In Section 3, we study the distribution and the popularity of descents on Catalan words avoiding each pattern of length three.

2 The sets \mathcal{C} and $\mathcal{C}(p)$ for $p \in \{00, 01, 10\}$

Here we consider both unrestricted Catalan words and those avoiding a length two pattern. We denote by $C(x, y)$ the bivariate generating function where the coefficient of $x^n y^k$ of its series expansion is the number of length n Catalan words with k descents. When we restrict to Catalan words avoiding the pattern p , the corresponding generating function is denoted by $C_p(x, y)$.

Theorem 1. *We have*

$$C(x, y) = \frac{1 - 2x + 2xy - \sqrt{1 - 4x + 4x^2 - 4x^2y}}{2xy}.$$

Proof. Let $w = 0(w'+1)w''$ be the first return decomposition of a non-empty Catalan word w with $w', w'' \in \mathcal{C}$. If w' (resp. w'') is empty then the number $d(w)$ of descents in w is the same as that of w'' (resp. w'); otherwise, we have $d(w) = d(w') + d(w'') + 1$ since there is a descent between $w'+1$ and w'' . So, we obtain the functional equation $C(x, y) = 1 + xC(x, y) + x(C(x, y) - 1) + xy(C(x, y) - 1)^2$ which gives the desired result. \square

As expected, $C(x) = C(x, 1) = \frac{1 - \sqrt{1 - 4x}}{2x}$ is the generating function for the Catalan numbers, and $\frac{\partial C(x, y)}{\partial y}|_{y=1}$ is the generating function for the descent popularity on \mathcal{C} , and we have the next corollary.

Corollary 1. *The popularity of descents on the set \mathcal{C}_n is $\binom{2n-2}{n-3}$, and its generating function is $\frac{1-4x+2x^2-(1-2x)\sqrt{1-4x}}{2x\sqrt{1-4x}}$ (sequence [A002694](#) in [11]).*

Catalan words of odd lengths encompass a smaller size Catalan structure. This result is stated in the next corollary, see the bold entries in Table 1.

$k \setminus n$	1	2	3	4	5	6	7	8	9	10
0	1	2	4	8	16	32	64	128	256	512
1			1	6	24	80	240	672	1792	4608
2					2	20	120	560	2240	8064
3							5	70	560	3360
4									14	252
Σ	1	2	5	14	42	132	429	1430	4862	16796

Table 1: Number $c_{n,k}$ of length n Catalan words with k descents for $1 \leq n \leq 10$ and $0 \leq k \leq 4$.

Corollary 2. *Catalan words of length $2n+1$ with n descents are enumerated by the n th Catalan number $\frac{1}{n+1} \binom{2n}{n}$.*

Proof. Clearly, the maximal number of descents in a length n Catalan word is $\lfloor \frac{n-1}{2} \rfloor$. Let w be a Catalan word of length $2n+1$ with n descents. We necessarily have $w = 0(w'+1)w''$ with $w', w'' \neq \epsilon$, $d(w') = \lfloor \frac{|w'|-1}{2} \rfloor$, $d(w'') = \lfloor \frac{|w''|-1}{2} \rfloor$ and $d(w) = d(w') + d(w'') + 1$. Since the length of w is odd, $|w'|$ and $|w''|$ have the same parity. If $|w'|$ and $|w''|$ are both even, then $d(w) = \frac{|w'|-2}{2} + \frac{|w''|-2}{2} + 1 = \frac{|w'|+|w''|-2}{2} < \lfloor \frac{(|w'|+|w''|+1)-1}{2} \rfloor = \lfloor \frac{n-1}{2} \rfloor$ which gives a contradiction. So, $|w'|$ and $|w''|$ are both odd, and we have $d(w) = \frac{|w'|-1}{2} + \frac{|w''|-1}{2} + 1 = \lfloor \frac{(|w'|+|w''|+1)-1}{2} \rfloor = \lfloor \frac{n-1}{2} \rfloor$. Thus the generating function $A(x)$ where the coefficient of x^n is the number of Catalan words of length $2n+1$ with n descents satisfies $A(x) = 1 + xA(x)^2$ which is the generating function for the Catalan numbers. \square

There are three patterns of length two, namely 00, 01 and 10, and Catalan words avoiding such a pattern do not have descents, thus the corresponding bivariate generating functions collapse into one variable ones.

Theorem 2. *For $p \in \{00, 01\}$, we have $C_p(x, y) = \frac{1}{1-x}$.*

Proof. If $p = 00$ (resp. $p = 01$) then $012\dots n-1$ (resp. $0\dots 0$) is the unique non-empty Catalan word of length n avoiding p , and the statement follows. \square

Theorem 3. *We have $C_{10}(x, y) = \frac{1-x}{1-2x}$, which is the generating function for the sequence 2^{n-1} (sequence [A011782](#) in [11]).*

Proof. A non-empty Catalan word avoiding the pattern 10 is of the form $0^k(w' + 1)$ for $k \geq 1$, and with $w' \in \mathcal{C}(10)$. So, we have the functional equation $C_{10}(x) = 1 + \frac{x}{1-x}C_{10}(x)$, which gives $C_{10}(x) = \frac{1-x}{1-2x}$. \square

3 The sets $\mathcal{C}(p)$ for a length three pattern p

Here we turn our attention to patterns of length three. There are thirteen such patterns, and we give the distribution and the popularity of descents on Catalan words avoiding each of them. Some of the obtained results are summarized in Tables 2 and 3.

Theorem 4. For $p \in \{012, 001\}$, we have

$$C_p(x, y) = \frac{1 - x + x^2 - x^2y}{1 - 2x + x^2 - x^2y}.$$

Proof. A non-empty word $w \in \mathcal{C}(012)$ has its first return decomposition $w = 01^k w''$ where $k \geq 0$ and $w'' \in \mathcal{C}(012)$. If $k = 0$ or $w'' = \epsilon$, then the number of descents in w is the same as that of w'' ; otherwise, we have $d(w) = d(w'') + 1$ (there is a descent between 1^k and w''). So, we obtain the functional equation $C_{012}(x, y) = 1 + xC_{012}(x, y) + \frac{x^2}{1-x} + \frac{x^2}{1-x}y(C_{012}(x, y) - 1)$ which gives the desired result.

A non-empty word $w \in \mathcal{C}(001)$ has the form $w = 0(w' + 1)0^k$ where $w' \in \mathcal{C}(001)$ and $k \geq 0$. If $k = 0$ or $w' = \epsilon$, then the number of descents in w is the same as that of w' ; otherwise, we have $d(w) = d(w') + 1$. So, we obtain the functional equation $C_{001}(x, y) = 1 + x(C_{001}(x, y) - 1) + \frac{x}{1-x} + \frac{x^2}{1-x}y(C_{001}(x, y) - 1)$ which gives the desired result. \square

Considering the previous theorem and the coefficient of x^n in $C_p(x, 1) = \frac{1-x}{1-2x}$ and in $\frac{\partial C_p(x, y)}{\partial y} \Big|_{y=1} = \frac{x^3}{(1-2x)^2}$, we obtain the next corollary.

Corollary 3. For $p \in \{012, 001\}$, we have $|\mathcal{C}_n(p)| = 2^{n-1}$, and the popularity of descents on the set $\mathcal{C}_n(p)$ is $(n-2) \cdot 2^{n-3}$ (sequence [A001787](#) in [11]).

As in the case of length two patterns, a Catalan word avoiding 010 does not have descents, and we have the next theorem.

Theorem 5. If $p = 010$, then $C_p(x, y) = \frac{1-x}{1-2x}$ which is the generating function for the sequence 2^{n-1} (sequence [A011782](#) in [11]).

Proof. A non-empty word $w \in \mathcal{C}(010)$ can be written either as $w = 0w'$ with $w' \in \mathcal{C}(10)$, or as $w = 0(w' + 1)$ with $w' \in \mathcal{C}(010) \setminus \{\epsilon\}$. So, we deduce $C_{010}(x) = 1 + xC_{10}(x) + x(C_{010}(x) - 1)$, and the statement holds. \square

Theorem 6. *For $p = 021$, we have*

$$C_p(x, y) = \frac{1 - 4x + 6x^2 - x^2y - 4x^3 + 3x^3y + x^4 - x^4y}{(1-x)(1-2x)(1-2x+x^2-x^2y)}.$$

Proof. Let w be a non-empty word in $\mathcal{C}(021)$, and let $0(w' + 1)w''$ its first return decomposition with $w', w'' \in \mathcal{C}(021)$. Note that w' belongs to $\mathcal{C}(10)$. We distinguish two cases: (1) w' does not contain 1, and (2) otherwise. In the case (1), $w' \in \mathcal{C}(01)$ (i.e., $w' = 0^k$ for some $k \geq 0$), and $w'' \in \mathcal{C}(021)$. If $w' = \epsilon$ (resp. $w'' = \epsilon$), then the number of descents in w is the same as that of w'' (resp. w'); otherwise, we have $d(w) = d(w') + d(w'') + 1$. So, this case contributes to $C_p(x, y)$ with $x C_{01}(x, y) + x(C_{021}(x, y) - 1) + xy(C_{01}(x, y) - 1)(C_{021}(x, y) - 1)$.

In the case (2), $w' \in \mathcal{C}(10) \setminus \mathcal{C}(01)$ and $w'' \in \mathcal{C}(01)$. If $w'' = \epsilon$ then w and w' have the same number of descents; otherwise, we have $d(w) = d(w') + d(w'') + 1$. So, this case contributes to $C_p(x, y)$ with $x(C_{10}(x, y) - C_{01}(x, y)) + xy(C_{10}(x, y) - C_{01}(x, y))(C_{01}(x, y) - 1)$.

Taking into account these two disjoint cases, and adding the empty word, we deduce the functional equation $C_{021}(x, y) = 1 + xC_{01}(x, y) + x(C_{021}(x, y) - 1) + xy(C_{01}(x, y) - 1)(C_{021}(x, y) - 1) + x(C_{10}(x, y) - C_{01}(x, y)) + xy(C_{10}(x, y) - C_{01}(x, y))(C_{01}(x, y) - 1)$, which after calculation gives the result. \square

Corollary 4. *For $p = 021$, we have $C_p(x) = \frac{1-4x+5x^2-x^3}{(1-2x)^2(1-x)}$ which is the generating function for the sequence $(n-1) \cdot 2^{n-2} + 1$ (sequence [A005183](#) in [11]). The popularity of descents on the set $\mathcal{C}_n(p)$ is $(n+1)(n-2) \cdot 2^{n-5}$ with the generating function $\frac{x^3(1-x)}{(1-2x)^3}$ (sequence [A001793](#) in [11]).*

Theorem 7. *For $p \in \{102, 201\}$, we have*

$$C_p(x, y) = \frac{1 - 3x + 3x^2 - 2x^2y - x^3 + x^3y}{(1-x)(1-3x+2x^2-2x^2y)}.$$

Proof. Let w be a non-empty word in $\mathcal{C}(102)$, and let $0(w' + 1)w''$ its first return decomposition with $w', w'' \in \mathcal{C}(102)$. If w' is empty, then $w = 0w''$ for some $w'' \in \mathcal{C}(102)$ and we have $d(w) = d(w'')$. If w'' is empty, then $w = 0(w' + 1)$ for some $w' \in \mathcal{C}(102)$ and we have $d(w) = d(w')$. If w' and w'' are both non-empty, then $w' \in \mathcal{C}(102) \setminus \{\epsilon\}$ and $w'' \in \mathcal{C}(012) \setminus \{\epsilon\}$. We deduce the functional equation $C_{102}(x, y) = 1 + xC_{102}(x, y) + x(C_{102}(x, y) -$

$1) + xy(C_{102}(x, y) - 1)(C_{012}(x, y) - 1)$. Finally, by Theorem 4 we obtain the desired result.

Let w be a non-empty word in $\mathcal{C}(201)$, and let $0(w' + 1)w''$ its first return decomposition with $w', w'' \in \mathcal{C}(201)$. If w' is empty, then $w = 0w''$ for some $w'' \in \mathcal{C}(201)$ and we have $d(w) = d(w'')$. If w'' is empty, then $w = 0(w' + 1)$ for some $w' \in \mathcal{C}(201)$ and we have $d(w) = d(w')$. If w' and w'' are both non-empty, then $d(w) = d(w') + d(w'') + 1$ and we distinguish two cases: (1) w' does not contain 1, and (2) otherwise. In the case (1), we have $w' \in \mathcal{C}(01) \setminus \{\epsilon\}$ and $w'' \in \mathcal{C}(201) \setminus \{\epsilon\}$; in the case (2), w' contains 1 and $w' \in \mathcal{C}(201) \setminus \mathcal{C}(01)$ and $w'' \in \mathcal{C}(01) \setminus \{\epsilon\}$. Combining the previous cases, the functional equation becomes $C_{201}(x, y) = 1 + xC_{201}(x, y) + x(C_{201}(x, y) - 1) + xy(C_{01}(x, y) - 1)(C_{201}(x, y) - 1) + xy(C_{201}(x, y) - C_{01}(x, y))(C_{01}(x, y) - 1)$, which gives the desired result. \square

Corollary 5. For $p \in \{102, 201\}$, we have $C_p(x) = \frac{1-3x+x^2}{(1-x)(1-3x)}$ which is the generating function of the sequence $\frac{3^{n-1}+1}{2}$ (sequence [A007051](#) in [11]). The popularity of descents on the set $\mathcal{C}_n(p)$ is $(n-2) \cdot 3^{n-3}$ with the generating function $\frac{x^3}{(1-3x)^2}$ (sequence [A027471](#) in [11]).

Theorem 8. For $p \in \{120, 101\}$, we have

$$C_p(x, y) = \frac{1 - 2x + x^2 - x^2y}{1 - 3x + 2x^2 - x^2y}.$$

Proof. Let w be a non-empty word in $\mathcal{C}(120)$, and let $0(w' + 1)w''$ be its first return decomposition where $w', w'' \in \mathcal{C}(120)$. If w'' is empty, then $w = 0(w' + 1)$ for some $w' \in \mathcal{C}(120)$ and we have $d(w) = d(w')$; if w' is empty, then $w = 0w''$ for some $w'' \in \mathcal{C}(120)$ and we have $d(w) = d(w'')$; if w' and w'' are not empty, then $w' \in \mathcal{C}(01) \setminus \{\epsilon\}$, $w'' \in \mathcal{C}(120) \setminus \{\epsilon\}$ and $d(w) = d(w') + d(w'') + 1$. We deduce the functional equation $C_{120}(x, y) = 1 + xC_{120}(x, y) + x(C_{120}(x, y) - 1) + xy(C_{01}(x, y) - 1)(C_{120}(x, y) - 1)$ which gives the result.

Let w be a non-empty word in $\mathcal{C}(101)$, and let $0(w' + 1)w''$ be its first return decomposition where $w', w'' \in \mathcal{C}(101)$. If w' is empty, then $w = 0w''$ for some $w'' \in \mathcal{C}(101)$ and $d(w) = d(w'')$; if w'' is empty, then $w = 0(w' + 1)$ for some $w' \in \mathcal{C}(101)$ and $d(w) = d(w')$; if w' and w'' are not empty, then $w' \in \mathcal{C}(101) \setminus \{\epsilon\}$ and $w'' \in \mathcal{C}(01) \setminus \{\epsilon\}$ and $d(w) = d(w') + d(w'') + 1$. We deduce the functional equation $C_{101}(x, y) = 1 + xC_{101}(x, y) + x(C_{101}(x, y) - 1) + xy(C_{101}(x, y) - 1)(C_{01}(x, y) - 1)$ which gives the result. \square

Corollary 6. For $p \in \{120, 101\}$, we have $C_p(x) = \frac{1-2x}{1-3x+x^2}$ and the coefficient of x^n in its series expansion is the $(2n-1)$ th term of the Fibonacci sequence (see [A001519](#) in [11]). The popularity of descents on the set $\mathcal{C}_n(p)$ is given by $\sum_{k=1}^{n-2} k \cdot \binom{n+k-2}{2k}$ which is the coefficient of x^n in the series expansion of $\frac{x^3(1-x)}{(1-3x+x^2)^2}$ (sequence [A001870](#) in [11]).

Theorem 9. For $p = 011$, we have

$$C_p(x, y) = \frac{1 - 2x + 2x^2 - x^3 + x^3y}{(1-x)^3}.$$

Proof. Let w be a non-empty word in $\mathcal{C}(011)$, and let $0(w'+1)w''$ its first return decomposition where $w', w'' \in \mathcal{C}(011)$. If w' (resp. w'') is empty, then we have $d(w) = d(w'')$ (resp. $d(w) = d(w')$); if w' and w'' are non-empty, then $w' \in \mathcal{C}(00) \setminus \{\epsilon\}$ and $w'' \in \mathcal{C}(01) \setminus \{\epsilon\}$. We deduce the functional equation $C_{011}(x, y) = 1 + xC_{011}(x, y) + x(C_{00}(x, y) - 1) + xy(C_{00}(x, y) - 1)(C_{01}(x, y) - 1)$ which gives the result. \square

Corollary 7. For $p = 011$, we have $C_p(x) = \frac{1-2x+2x^2}{(1-x)^3}$ and the coefficient of x^n in its series expansion is $1 + \binom{n}{2}$ (sequence [A000124](#) in [11]). The popularity of descents on the set $\mathcal{C}_n(p)$ is given by $\frac{(n-1)(n-2)}{2}$ which is the coefficient of x^n in the series expansion of $\frac{x^3}{(1-x)^3}$ (sequence [A000217](#) in [11]).

Theorem 10. For $p = 000$, we have

$$C_p(x, y) = \frac{1 - x^2 - x^2y}{1 - x - 2x^2 - x^2y + x^3 + x^4 - x^4y}.$$

Proof. Let w be a non-empty word in $\mathcal{C}(000)$, and let $0(w'+1)w''$ its first return decomposition where $w', w'' \in \mathcal{C}(000)$. We distinguish two cases: (1) w'' is empty, and (2) otherwise.

In the case (1), we have $w = 0(w'+1)$ for some $w' \in \mathcal{C}(000)$ and $d(w) = d(w')$. So, the generating function $A(x, y)$ for the Catalan words in this case is $A(x, y) = xC_{000}(x, y)$.

In the case (2), we set $w'' = 0(w''' + 1)$ for some $w''' \in \mathcal{C}(000)$ and we have $w = 0(w'+1)0(w''' + 1)$. We distinguish three sub-cases: (2.a) w' is empty, (2.b) w' is non-empty and w''' is empty, and (2.c) w' and w''' are both non-empty.

In the case (2.a), we have $w = 00(w''' + 1)$ with $w''' \in \mathcal{C}(000)$. So, the generating function for the Catalan words belonging to this case is $B_a(x, y) = x^2C_{000}(x, y)$.

In the case (2.b), we have $w = 0(w' + 1)0$ with $w' \in \mathcal{C}(000) \setminus \{\epsilon\}$. So, the generating function for the corresponding Catalan words is $B_b(x, y) = x^2y(C_{000}(x, y) - 1)$.

In the case (2.c), we have $w = 0(w' + 1)0(w''' + 1)$ where w' and w''' are non-empty Catalan words such that $w'w'''$ is a Catalan word lying in the case (2). If $w' = 0$, then $d(w'w''') = d(w''') = d(w) - 1$; if $w' \neq 0$, then $d(w'w''') = d(w') + d(w''') + 1 = d(w)$. So, the generating function for the corresponding Catalan words is $B_c(x, y) = x^2yB_a(x, y) + x^2(B_b(x, y) + B_c(x, y))$.

Considering $C_{000}(x, y) = 1 + A(x, y) + B_a(x, y) + B_b(x, y) + B_c(x, y)$, the obtained functional equations give the result. \square

Corollary 8. For $p = 000$, we have $C_p(x) = \frac{1-2x^2}{1-x-3x^2+x^3}$ and the generating function for the popularity of descents in the sets $\mathcal{C}_n(p)$, $n \geq 0$, is

$$\frac{x^3(1-x)(1+2x)(1+x)}{(1-x-3x^2+x^3)^2}.$$

Note that the sequences defined by the two generating functions in Corollary 8 do not appear in [11].

Theorem 11. For $p = 100$, we have

$$C_p(x, y) = \frac{1 - 2x - x^2y + x^3}{1 - 3x + x^2 - x^2y + 2x^3}.$$

Proof. For $k \geq 1$, we define $\mathcal{A}_k \subset \mathcal{C}(100)$ as the set of Catalan words avoiding 100 with exactly k zeros, and let $A_k(x, y)$ be the generating function for \mathcal{A}_k .

A Catalan word $w \in \mathcal{A}_1$ is of the form $w = 0(w' + 1)$ with $w' \in \mathcal{C}(100)$. Since we have $d(w) = d(w')$, the generating function $A_1(x, y)$ for these words satisfies $A_1(x, y) = xC_{100}(x, y)$.

A Catalan word $w \in \mathcal{A}_k$, $k \geq 3$, is of the form $w = 0^{k-2}w'$ with $w' \in \mathcal{A}_2$. Since we have $d(w) = d(w')$, the generating function $A_k(x, y)$ for these words satisfies $A_k(x, y) = x^{k-2}A_2(x, y)$.

A Catalan word $w \in \mathcal{A}_2$ has one of the three following forms:

(1) $w = 00(w' + 1)$ with $w' \in \mathcal{C}(100)$; we have $d(w) = d(w')$, and the generating function for these Catalan words is $x^2C_{100}(x, y)$.

(2) $w = 0(w' + 1)0$ with $w' \in \mathcal{C}(100) \setminus \{\epsilon\}$; we have $d(w) = d(w') + 1$, and the generating function for these Catalan words is $x^2y(C_{100}(x, y) - 1)$.

(3) $w = 0(w' + 1)0(w'' + 1)$ where w' and w'' are non-empty and $w'w'' \in \mathcal{A}_k$ for some $k \geq 2$ (i.e., $w'w'' = 0^{k-2}0(u+1)0(v+1)$ with $0(u+1)0(v+1) \in \mathcal{A}_2$). So, there are $(k-1)$ possible choices for w' , namely $0, 0^2, \dots, 0^{k-2}$, and $0^{k-2}0(u+1)$. If $w' = 0, 0^2, \dots, 0^{k-2}$, then $d(w) = d(0(u+1)0(v+1)) +$

1; if $w' = 0^{k-2}0(u+1)$ and $u \neq \epsilon$, then $d(w) = d(0(u+1)0(v+1))$; if $w' = 0^{k-2}0(u+1)$ and $u = \epsilon$, then $d(w) = d(0(u+1)0(v+1)) + 1$. So, the generating function for these words is $x^2yA_2(x, y) \sum_{k \geq 2} (k-2)x^{k-2} + x^2(A_2(x, y) - x^2C_{100}(x, y)) \sum_{k \geq 2} x^{k-2} + x^4yC_{100}(x, y) \sum_{k \geq 2} x^{k-2}$, which is $\frac{x^3y}{(1-x)^2}A_2(x, y) + \frac{x^2}{1-x}A_2(x, y) + \frac{x^4y-x^2}{1-x}C_{100}(x, y)$.

Taking into account all previous cases, we obtain the following functional equations:

(i) $A_1(x, y) = xC_{100}(x, y)$,

(ii) $A_2(x, y) = x^2C_{100}(x, y) + x^2y(C_{100}(x, y) - 1) + \frac{x^3y}{(1-x)^2}A_2(x, y) + \frac{x^2}{1-x}A_2(x, y) + \frac{x^4y-x^2}{1-x}C_{100}(x, y)$,

(iii) $A_k(x, y) = x^{k-2}A_2(x, y)$ for $k \geq 3$,

(iv) $C_{100}(x, y) = 1 + \sum_{k \geq 1} A_k(x, y)$.

A simple calculation gives the desired result. \square

Corollary 9. For $p = 100$, we have $C_p(x) = \frac{1-2x-x^2+x^3}{1-3x+2x^3}$, which is the generating function for the sequence $\lceil \frac{(1+\sqrt{3})^{n+1}}{12} \rceil$ (see [A057960](#) in [11]), and the generating function for the popularity of descents in the sets $\mathcal{C}_n(p)$, $n \geq 0$, is

$$\frac{x^3(1-x-x^2)}{(1-3x+2x^3)^2}.$$

Theorem 12. For $p = 110$, we have

$$C_p(x) = \frac{1-3x+2x^2+x^3-x^4+x^4y}{(1-x)(1-3x+x^2+2x^3-x^3y)}.$$

Proof. Let w be a non-empty word in $\mathcal{C}(110)$, and let $0(w'+1)w''$ its first return decomposition where $w', w'' \in \mathcal{C}(110)$.

Then, w has one of the following forms:

- $w = 0(w'+1)$ where $w' \in \mathcal{C}(110)$; the generating function for these words is $xC_{110}(x, y)$.
- $w = 0w'$ where $w' \in \mathcal{C}(110) \setminus \{\epsilon\}$; the generating function for these words is $x(C_{110}(x, y) - 1)$.
- $w = 0(w'+1)w''$ with $w' \in \mathcal{C}(00) \setminus \{\epsilon\}$ and $w'' \in \mathcal{C}(10) \setminus \{\epsilon\}$; the generating function for these words is $xy(C_{00}(x, y) - 1)(C_{10}(x, y) - 1)$.

- The last form is $w = 0(w'+1)w''$ where $w' \in \mathcal{C}(00) \setminus \{\epsilon\}$ and $w'' \notin \mathcal{C}(10)$. So, we have $w = 012 \dots k0^{a_0}1^{a_1} \dots (k-1)^{a_k}(w''' + k - 1)$ where $k \geq 1$, $a_i \geq 1$ for $0 \leq i \leq k$, and $w''' \in \mathcal{C}(110) \setminus \mathcal{C}(10)$; the generating function for these words is $y \sum_{k \geq 1} \frac{x^{2k+1}}{(1-x)^k} (C_{110}(x, y) - C_{10}(x, y))$.

Combining these different cases, we deduce the functional equation:

$$C_{110}(x, y) = 1 + xC_{110}(x, y) + x(C_{110}(x, y) - 1) + xy(C_{00}(x, y) - 1)(C_{10}(x, y) - 1) + y \sum_{k \geq 1} \frac{x^{2k+1}}{(1-x)^k} (C_{110}(x, y) - C_{10}(x, y)).$$

Considering Theorems 3 and 2, the result follows. \square

Corollary 10. For $p = 110$, we have $C_p(x) = \frac{1-3x+2x^2+x^3}{(1-x)^2(1-2x-x^2)}$ and the generating function for the popularity of descents in the sets $\mathcal{C}_n(p)$, $n \geq 0$, is

$$\frac{x^3(1-x-x^2)^2}{(1-x)^3(1-2x-x^2)^2}.$$

Theorem 13. For $p = 210$, we have

$$C_p(x) = \frac{1-5x+8x^2-x^2y-4x^3+3x^3y-x^4y}{(1-2x)(1-4x+4x^2-x^2y+x^3y)}.$$

Proof. Let w be a non-empty word in $\mathcal{C}(210)$, and let $0(w'+1)w''$ be its first return decomposition where $w', w'' \in \mathcal{C}(210)$.

Then, w has one of the following forms:

- $w = 0(w'+1)$ where $w' \in \mathcal{C}(210)$; the generating function for these words is $x C_{210}(x, y)$.
- $w = 0w''$ where $w'' \in \mathcal{C}(210) \setminus \{\epsilon\}$; the generating function for these words is $x(C_{210}(x, y) - 1)$.
- $w = 0(w'+1)w''$ where $w' \in \mathcal{C}(01) \setminus \{\epsilon\}$ and $w'' \in \mathcal{C}(210) \setminus \{\epsilon\}$; the generating function for these sets is $xy(C_{01}(x, y) - 1)(C_{210}(x, y) - 1)$.
- $w = 01^{a_1}2^{a_2} \dots k^{a_k}w''$ where $k \geq 2$, $a_i \geq 1$ for $1 \leq i \leq k$, and $w'' \in \mathcal{C}(10) \setminus \{\epsilon\}$; the generating function for these words is $y(C_{10}(x, y) - 1) \sum_{k \geq 2} \frac{x^{k+1}}{(1-x)^k}$.
- $w = 01^{a_1}2^{a_2} \dots k^{a_k}0^{b_0}1^{b_1} \dots (k-2)^{b_{k-2}}(w'' + k - 2)$ where $k \geq 2$, $a_i \geq 1$ for $1 \leq i \leq k$, $b_i \geq 1$ for $1 \leq i \leq k-2$, and $w'' \in \mathcal{C}(210) \setminus \mathcal{C}(10)$; the generating function for these words is $y(C_{210}(x, y) - C_{10}(x, y)) \sum_{k \geq 2} \frac{x^{k+1}}{(1-x)^k} \frac{x^{k-1}}{(1-x)^{k-1}}$.

Pattern p	Sequence $ \mathcal{C}_n(p) $	Generating function	OEIS [11]
012, 001, 010	2^{n-1}	$\frac{1-x}{1-2x}$	A011782
021	$(n-1) \cdot 2^{n-2} + 1$	$\frac{1-4x+5x^2-x^3}{(1-x)(1-2x)^2}$	A005183
102, 201	$\frac{3^{n-1}+1}{2}$	$\frac{1-3x+x^2}{(1-x)(1-3x)}$	A007051
120, 101	F_{2n-1}	$\frac{1-2x}{1-3x+x^2}$	A001519
011	$\frac{n(n-1)}{2} + 1$	$\frac{1-2x+2x^2}{(1-x)^3}$	A000124
000		$\frac{1-2x^2}{1-x-3x^2+x^3}$	
100	$\lceil \frac{(1+\sqrt{3})^{n+1}}{12} \rceil$	$\frac{1-2x-x^2+x^3}{1-3x+2x^3}$	A057960
110	$\frac{1}{2} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n+1}{2k+1} 2^k - \frac{n-1}{2}$	$\frac{1-3x+2x^2+x^3}{(1-x)^2(1-2x-x^2)}$	
210		$\frac{1-5x+7x^2-x^3-x^4}{(1-2x)(1-4x+3x^2+x^3)}$	

Table 2: Catalan words avoiding a pattern of length three.

Combining these different cases, we deduce the functional equation:

$$C_{210}(x, y) = 1 + xC_{210}(x, y) + x(C_{210}(x, y) - 1) + xy(C_{01}(x, y) - 1)(C_{210}(x, y) - 1) + y(C_{10}(x, y) - 1) \sum_{k \geq 2} \frac{x^{k+1}}{(1-x)^k} + y(C_{210}(x, y) - C_{10}(x, y)) \sum_{k \geq 2} \frac{x^{k+1}}{(1-x)^k} \frac{x^{k-1}}{(1-x)^{k-1}}.$$

Finally, considering Theorem 3 the desired result follows. \square

Corollary 11. For $p = 210$, we have $C_p(x) = \frac{1-5x+7x^2-x^3-x^4}{(1-2x)(1-4x+3x^2+x^3)}$ and the generating function for the popularity of descents in the set $\mathcal{C}_n(p)$, $n \geq 0$, is

$$\frac{x^3(1-2x)}{(1-4x+3x^2+x^3)^2}.$$

Pattern p	Popularity of descents on $\mathcal{C}_n(p)$	Generating function	OEIS [11]
012, 001	$(n-2) \cdot 2^{n-3}$	$\frac{x^3}{(1-2x)^2}$	A001787
010	0	0	
021	$(n+1)(n-2) \cdot 2^{n-5}$	$\frac{x^3(1-x)}{(1-2x)^3}$	A001793
102, 201	$(n-2) \cdot 3^{n-3}$	$\frac{x^3}{(1-3x)^2}$	A027471
120, 101	$\sum_{k=1}^{n-2} k \cdot \binom{n+k-2}{2k}$	$\frac{x^3(1-x)}{(1-3x+x^2)^2}$	A001870
011	$\frac{(n-1)(n-2)}{2}$	$\frac{x^3}{(1-x)^3}$	A000217
000		$\frac{x^3(1-x)(1+2x)(1+x)}{(1-x-3x^2+x^3)^2}$	
100		$\frac{x^3(1-x-x^2)}{(1-3x+2x^3)^2}$	
110		$\frac{x^3(1-x-x^2)^2}{(1-x)^3(1-2x-x^2)^2}$	
210		$\frac{x^3(1-2x)}{(1-4x+3x^2+x^3)^2}$	

Table 3: Popularity of descents on Catalan words avoiding a pattern of length three.

4 Final remarks

At the time of writing this paper, the enumerating sequences $(|\mathcal{C}_n(p)|)_{n \geq 0}$, for $p \in \{000, 110, 210\}$, are not recorded in [11], and it will be interesting to explore potential connections of these sequences with other known ones.

According to Theorem 4, for any $k \geq 0$, the set of fixed length Catalan words with k descents avoiding $p = 001$ is equinumerous with those avoiding $q = 012$, and a natural question that arises is to find a constructive bijection between the two sets; and similarly for $(p, q) = (102, 201)$, see Theorem 7, and for $(p, q) = (101, 120)$, see Theorem 8. In the same vein, some of the enumerating sequences obtained in this paper count classical combinatorial classes (see Tables 2 and 3) and these results deserve bijective proofs.

Finally, our initiating study on pattern avoidance on Catalan words can naturally be extended to patterns of length more than three, vincular patterns and/or multiple pattern avoidance. For example, some of the patterns we considered here hide larger length patterns (for instance, an occurrence of 210 in a Catalan word is a part of an occurrence of 01210), and some of our results can be restated in this light.

References

- [1] M. Albert, C. Homberger, and J. Pantone. Equipopularity classes in the separable permutations. *The Electronic J. of Comb.*, 22(2):P2.2, 2015. (electronic).
- [2] J.-L. Baril, S. Kirgizov, and V. Vajnovszki. Patterns in treeshelves. *Discrete Mathematics*, 340(12):2946–2954, 2017.
- [3] F. Bergeron, N. Bergeron, R.B. Howlett, and D.E. Taylor. A decomposition of the descent algebra of a finite Coxeter group. *J. Algebraic Combin.*, 1:23–44, 1992.
- [4] M. Bóna. Surprising symmetries in objects counted by Catalan numbers. *The Electronic J. of Comb.*, 19(1):P62, 2012. (electronic).
- [5] E. Deutsch. Dyck path enumeration. *Discrete Math.*, 204:167–202, 1999.
- [6] I. Gessel and G. Viennot. Binomial determinants, paths, and hook length formulae. *Adv. Math.*, 58:300–321, 1985.

- [7] C. Homberger. Expected patterns in permutation classe. *The Electronic J. of Comb.*, 19(3):P43, 2012. (electronic).
- [8] E. Mäkinen. Left distance binary tree representations. *BIT Numerical Mathematics*, 27(2):163–169, 1987.
- [9] T. Mansour and V. Vajnovszki. Efficient generation of restricted growth words. *Information Processing Letters*, 113:613–616, 2013.
- [10] K. Rudolph. Pattern popularity in 132-avoiding permutations. *The Electronic J. of Comb.*, 20(1):P8, 2013. (electronic).
- [11] N.J.A. Sloane. The on-line encyclopedia of integer sequences. Available electronically at <http://oeis.org>.
- [12] R.P. Stanley. *Enumerative Combinatorics*, volume 2. Cambridge University Press, 1999.