

# Forests and pattern avoiding permutations modulo pure descents

Jean-Luc Baril, Sergey Kirgizov and Armen Petrossian

LE2I, Université de Bourgogne, B.P. 47 870, 21078 Dijon, France  
{barjl, sergey.kirgizov, armen.petrossian}@u-bourgogne.fr

## Abstract

We investigate an equivalence relation on permutations based on the pure descent statistic. Generating functions are given for the number of equivalence classes for some restricted sets of permutations avoiding at most one pattern of length three. As a byproduct, we exhibit a permutation set in one-to-one correspondence with forests of ordered binary trees, which provides a new combinatorial class enumerated by the single-source directed animals on the square lattice. Furthermore, bivariate generating functions for these sets are given according to various statistics.

**Keywords:** permutation; equivalence class; pure descent; pattern; Catalan and Motzkin numbers; forest; directed animals.

## 1 Introduction, definitions and notations

Many statistics on permutations have been studied for many years, but two of them appear more frequently in the literature: the number of descents and the number of excedances. These two statistics have been introduced by MacMahon [12] and are closely related since they have the same distribution. However, many more articles deal with the descents which have links with other fields as the Coxeter group [6] or the theory of lattice path [9].

Recently [3, 4], two equivalence relations on permutations based on the excedance and descent statistics have been introduced. The main results of these works consist in giving generating functions for the number of equivalence classes for several restricted

sets of permutations such as involutions, cycles, derangements, permutations avoiding at most one pattern of length three. So, it becomes natural to conduct a similar study for equivalence relations based on other statistics. It is precisely the aim of this paper which investigates an equivalence relation based on the pure descent statistic, first introduced in [5] and formally defined below. Moreover, we show how equivalence classes are in one-to-one correspondence with certain forests of ordered trees, providing some links between several statistics on these sets. As a consequence, we exhibit a new set of pattern avoiding permutations with the same cardinality as the set of single-source directed animals on the square lattice (see Barucci and al. [2], Bousquet-Mélou [8] for two studies concerning directed animals, forests and pattern avoiding permutations).

Now, we present some basic definitions and notations. Let  $S_n$  be the set of permutations of length  $n$ , *i.e.*, all one-to-one correspondences from  $[n] = \{1, 2, \dots, n\}$  into itself. The *one-line notation* of a permutation  $\pi \in S_n$  is  $\pi_1\pi_2 \cdots \pi_n$  where  $\pi_i = \pi(i)$  for  $i \in [n]$ . The *graphical representation* of  $\pi \in S_n$  is the set of points in the plane at coordinates  $(i, \pi_i)$  for  $i \in [n]$  (see Figure 1).

Let  $\pi$  be a permutation in  $S_n$ . A *descent* of  $\pi$  is an integer  $i \in [n - 1]$  such that  $\pi_i > \pi_{i+1}$ . Whenever there does not exist  $j < i$  such that  $\pi_{i+1} < \pi_j < \pi_i$ , we call it a *pure descent*. Let  $D(\pi)$  be the set of pure descents in  $\pi$ , and  $DD(\pi)$  be the set of pairs  $(\pi_i, \pi_{i+1})$  for  $i \in D(\pi)$ . By abuse of language, such a pair will be also called a *pure descent*. For instance, if  $\pi = 1\ 4\ 2\ 7\ 5\ 3\ 8\ 6$  then  $D(\pi) = \{2, 4\}$  and  $DD(\pi) = \{(4, 2), (7, 5)\}$ . In [5, Theorem 1], the authors prove that the number of  $n$ -length permutations with  $k$  pure descents is given by the signless Stirling number of the first kind  $c(n, k + 1)$  where  $c(n, k)$  satisfies

$$c(n, k) = (n - 1) \cdot c(n - 1, k) + c(n - 1, k - 1)$$

with the initial conditions  $c(n, k) = 0$  if  $n \leq 0$  or  $k \leq 0$ , except  $c(0, 0) = 1$  ([19, 23] and the sequence [A132393](#) in the Sloane's on-line encyclopedia of integer sequences [18]).

We define on permutations of the same length the following equivalence relation based on the pure descent statistic:

$$\pi \sim \sigma \iff DD(\pi) = DD(\sigma).$$

The set of equivalence classes in  $S_n$  (resp. in a restricted set  $R \subset S_n$ ) is denoted  $S_n^\sim$  (resp.  $R^\sim$ ). For instance, the permutations  $\pi = 1\ 4\ 2\ 7\ 5\ 3\ 8\ 6$  and  $\sigma = 1\ 7\ 5\ 6\ 4\ 2\ 3\ 8$  belong to the same equivalence class (see Figure 1) because  $DD(\pi) = DD(\sigma) = \{(4, 2), (7, 5)\}$ , and  $S_3^\sim$  is constituted of the 5 classes  $\{123, 231\}$ ,  $\{132\}$ ,  $\{213\}$ ,  $\{321\}$  and  $\{312\}$ .




Figure 1: Two permutations  $\pi = 1\ 4\ 2\ 7\ 5\ 3\ 8\ 6$  and  $\sigma = 1\ 7\ 5\ 6\ 4\ 2\ 3\ 8$  in the same class with  $DD(\pi) = DD(\sigma) = \{(4, 2), (7, 5)\}$ .

A permutation  $\pi \in S_n$  *avoids* the pattern  $\tau \in S_k$  if and only if there is no sequence of indices  $1 \leq i_1 < i_2 < \dots < i_k \leq n$  such that  $\pi(i_1)\pi(i_2)\dots\pi(i_k)$  is order-isomorphic to  $\tau$  (see [17, 20]). We denote by  $S_n(\tau)$  the set of permutations of  $S_n$  avoiding the pattern  $\tau$ . For example, if  $\tau = 123$  then  $52143 \in S_5(\tau)$  while  $21534 \notin S_5(\tau)$ . Many classical sequences in combinatorics appear as the cardinality of pattern-avoiding permutation sets. A large number of these results were firstly obtained by West and Knuth [11, 16, 17, 20, 21, 22]. Also, we refer to the books of Kitaev [10], Mansour [14] and Bóna [7].

Later, Babson and Steingrímsson [1] defined *generalized patterns* (also called *vincular patterns*) where any pair of two adjacent values in the pattern may be underlined, which means that the corresponding values in the permutation must be adjacent. For instance, the pattern  $2\underline{31}$  occurs in the permutation 316452 two times as the subsequences 352 and 452, while the pattern 231 occurs four times.

Moreover, we will consider a *barred pattern*  $\bar{\tau}$ , *i.e.*, a permutation in  $S_k$  having a bar over one value (see [15] for instance). Let  $\tau$  be the permutation on  $[k]$  identical to  $\bar{\tau}$  but unbarred, and  $\hat{\tau}$  be the permutation on  $[k - 1]$  made up of the  $k - 1$  unbarred values of  $\bar{\tau}$  rewritten to be a permutation on  $[k - 1]$ . Then  $\pi \in S_n$  avoids the pattern  $\bar{\tau}$  if and only if each pattern  $\hat{\tau}$  in  $\pi$  can be expanded into a pattern  $\tau$  in  $\pi$  where the expanded value corresponds to the barred value in  $\bar{\tau}$ . For instance, the permutation 3241 does not avoid  $21\bar{3}$  since 41 cannot be expanded into a pattern 213, while 3124 avoids it.

In the following, we will consider permutations avoiding the generalized and barred pattern  $\underline{51}\bar{4}23$ , consisting in permutations where any pattern  $\underline{41}23$  can be expanded into a pattern  $\underline{51}423$ .

The main goal of this paper is to calculate the number of equivalence classes (modulo pure descents) for some subsets of permutations avoiding at most one pattern of length three. See Table 1 for an overview of our results.

The paper is organized as follows. In Section 2, we give a one-to-one correspondence

between  $S_n^\sim$  and the set of noncrossing partitions of  $[n]$ , proving that the cardinalities of  $S_n^\sim$  for  $n \geq 1$  are given by the Catalan numbers (see [A000108](#) in the On-line Encyclopedia of Integer Sequences [18]). For the case of permutations avoiding the pattern 231, we prove that any equivalence class contains only one permutation on which we construct a forest of ordered trees. Also, we prove that  $S_n(312)^\sim$  and  $S_n(321)^\sim$  are enumerated by  $2^{n-1}$  ([A011782](#) in [18]).

In Section 3, we describe a bijection between forests of ordered binary trees with  $n$  nodes and the set  $S_n(231, \underline{51\bar{4}23})$ , giving a new set of pattern avoiding permutations in bijection with the single-source directed animals on the square lattice (see Barucci and al. [2, 8]). Bivariate generating functions are given for these sets according to various statistics.

In Section 4, we investigate the equivalence relation on the set  $S_n(123)$  of permutation avoiding the pattern 123. We give a constructive bijection between forests of ordered binary trees and the classes in  $S_n(123)^\sim$ , proving that the cardinality of  $S_n(123)^\sim$  is also given by the sequence [A005773](#) that counts the single-source directed animals as above.

Pattern	Sequence	Sloane	$a_n, 1 \leq n \leq 9$
$\{\}, \{231\}$	Catalan	<a href="#">A000108</a>	1, 2, 5, 14, 42, 132, 429, 1430, 4862
$\{312\}, \{321\}$	$2^{n-1}$	<a href="#">A011782</a>	1, 2, 4, 8, 16, 32, 64, 128, 256
$\{231, \underline{51\bar{4}23}\}$	Directed animals	<a href="#">A005773</a>	1, 2, 5, 13, 35, 96, 267, 750, 2123
$\{123\}$	Directed animals	<a href="#">A005773</a>	1, 2, 5, 13, 35, 96, 267, 750, 2123

Table 1: Number of equivalence classes for some restricted sets of pattern avoiding permutations.

## 2 Enumeration of $S_n^\sim$ , $S_n(231)^\sim$ , $S_n(312)^\sim$ and $S_n(321)^\sim$

In this section, we provide the cardinality of  $S_n^\sim$ ,  $S_n(231)^\sim$ ,  $S_n(312)^\sim$  and  $S_n(321)^\sim$ . Note that if a permutation  $\pi$  avoids the pattern 231 then any descent of  $\pi$  is a pure descent.

A *partition*  $\Pi$  of  $[n]$  is a collection of non-empty pairwise disjoint subsets, called *blocks*, whose union is  $[n]$  (see [13]). The *standard form* of  $\Pi$  is  $\Pi_1/\Pi_2/\dots$ , where the blocks  $\Pi_i$  are arranged in increasing order of their smallest elements, and elements in a same block are in decreasing order. Let  $\mathcal{P}_n$  be the set of partitions of  $[n]$ , and

$\mathcal{NCP}_n \subset \mathcal{P}_n$  be the set of noncrossing partitions, *i.e.* all partitions  $\Pi$  where there do not exist four integers  $p < q < r < s$  such that  $p, r \in \Pi_i$  and  $q, s \in \Pi_j$  with  $i \neq j$ .

We associate to a permutation  $\pi \in S_n$  the unique partition  $\Pi$  defined as follows. Two elements  $x$  and  $y$ ,  $x > y$ , belong to the same block in  $\Pi$  if and only if there exist  $i$  and  $j$ ,  $i < j$ , such that the pairs  $(x = \pi_i, \pi_{i+1}), (\pi_{i+1}, \pi_{i+2}), \dots, (\pi_{j-1}, \pi_j = y)$  are pure descents in  $\pi$ . For instance, the two permutations in Figure 1 are associated to the same partition (in standard form)  $\Pi = 1/42/3/75/6/8$ . Moreover, two permutations in the same equivalence class (modulo pure descents) are associated to the same noncrossing partitions. Indeed, let us consider a block  $\Pi_i = \pi_a \pi_{a+1} \dots \pi_b$  with  $a < b$ ,  $\pi_a > \pi_{a+1} > \dots > \pi_b$ . Since  $\pi_a \pi_{a+1} \dots \pi_b$  is a subsequence of consecutive pure descents in  $\pi$ , there is no  $\pi_c$ ,  $c < a$ , such that  $\pi_c \in [\pi_b, \pi_a]$ . So, let us assume that there is  $c > b$  such that  $\pi_c \in [\pi_b, \pi_a]$ ; then, for the same argument, all elements in the same block as  $\pi_c$  are greater than  $\pi_b$  and lower than  $\pi_a$ , which implies that there is no  $\pi_d$  in the block of  $\pi_c$  such that  $\pi_d < \pi_b < \pi_c < \pi_a$  with  $a < b < c < d$ . Thus, the partition  $\Pi$  is noncrossing.

Conversely, any noncrossing partition  $\Pi$  of standard form  $\Pi = \Pi_1/\Pi_2/\dots/\Pi_k$ ,  $k \geq 1$ , is associated to the permutation  $\pi = \Pi_1\Pi_2\dots\Pi_k$  that avoids the pattern 231. Indeed, the noncrossing property forces all descents of  $\pi$  to be pure, implying that  $\pi$  does not contain any pattern 231. As the set  $\mathcal{NCP}_n$  (and also  $S_n(231)$ ) is enumerated by the  $n$ th Catalan number (see A000108, [18]), we obtain Theorem 1. As an immediate consequence, equivalence classes in  $S_n(231)^\sim$  are singletons, and the set  $S_n(231)$  is a set of representatives of  $S_n^\sim$ .

**Theorem 1** *The sets  $S_n^\sim$  (resp.  $S_n(231)^\sim$ ),  $n \geq 1$ , are enumerated by the Catalan numbers.*

As a byproduct of Theorem 1, we obtain the cardinalities of  $S_n(312)^\sim$  and  $S_n(321)^\sim$ . Since  $S_n(231)$  is a set of representatives of  $S_n^\sim$  (see Theorem 1), there is a unique  $\pi' \in S_n(231)$  equivalent to  $\pi \in S_n(312)$ , and  $\pi'$  is obtained from the noncrossing partition associated to  $\pi$  (in standard form)  $\Pi = \Pi_1/\Pi_2/\dots/\Pi_k$  by deleting all '/', *i.e.*,  $\pi' = \Pi_1\Pi_2\dots\Pi_k$ . Notice that for any permutation  $\pi \in S_n(312)$ , a pure descent in  $\pi$  is necessarily an adjacency, *i.e.*, a descent  $(\pi_i, \pi_{i+1})$  with  $\pi_{i+1} = \pi_i - 1$ . Then, any block  $\Pi_j$ ,  $1 \leq j \leq k$ , is an interval, which implies that  $\pi'$  avoids also the pattern 312. So, the set  $S_n(312)^\sim$  is in one-to-one correspondence with the set of  $S_n(231, 312)$  which induces Theorem 2 (see Simion and Schmidt [17]). Theorem 3 is obtained *mutatis mutandis*.

**Theorem 2** *The sets  $S_n(312)^\sim$ ,  $n \geq 1$ , are enumerated by the  $2^{n-1}$ .*

**Theorem 3** *The sets  $S_n(321)^\sim$ ,  $n \geq 1$ , are enumerated by the  $2^{n-1}$ .*

### 3 Forests and 231-avoiding permutations

In this section, we establish a constructive bijection between  $S_n(231)$  and the set  $\mathcal{F}_n$  of forests of ordered trees, *i.e.* collections of rooted trees in which children of each node are ordered and the total number of nodes is  $n$ . Taking advantage of the recursive definition of the forests, we exhibit a new set of permutations  $S_n(231, \underline{51\bar{4}23})$  having the same cardinality as the set of single-source directed animals on the square lattice (see [2, 8] and A005773, [18]). Moreover, we show how the bijection transports various statistics (see Table 2). As a byproduct, we provide several bivariate generating functions with respect to the length and these statistics for the two sets  $S_n(231)$  and  $S_n(231, \underline{51\bar{4}23})$  (see Table 3).

Let  $\pi$  be a permutation in  $S_n(231)$ . We construct a forest  $f_\pi \in \mathcal{F}_n$  as follows: we cross the graphical representation of  $\pi$  from left to right; if the point  $(i, \pi_i)$  is a left-to-right maximum (that is  $\pi_i > \pi_j$  for all  $j < i$ ), then it corresponds to the root of a new tree in  $f_\pi$ ; otherwise we add an edge between  $(i, \pi_i)$  and  $(j, \pi_j)$  where  $j$  is the rightmost  $j < i$  such that  $\pi_j > \pi_i$ . See Figure 2 for an example of this construction.

By construction, the map  $\phi : S_n(231) \rightarrow \mathcal{F}_n$  defined by  $\pi \mapsto f_\pi$  is injective. Since  $\mathcal{F}_n$  is enumerated by the  $n$ th Catalan number as  $S_n(231)$ , we deduce the bijectivity of  $\phi$ .




Figure 2: The permutation  $\pi = 8 4 1 2 3 6 5 7 9 13 11 10 12$  with its corresponding forest  $f_\pi$  and the noncrossing partition  $\Pi = 8 4 1/2/3/6 5/7/9/13 11 10/12$ .

Now we define some statistics on  $S_n(231)$  and  $\mathcal{F}_n$ , and we show how the map  $\phi$  establishes a correspondence between them.

For a permutation  $\pi \in S_n(231)$ , we define:

- des**( $\pi$ ) = number of descents (which is also the number of pure descents);
- ides**( $\pi$ ) = number of descents in  $\pi^{-1}$  (for  $\pi \in S_n(231)$ , we have **ides**( $\pi$ ) = **des**( $\pi$ ));
- adj**( $\pi$ ) = number of adjacencies, i.e. descent  $(\pi_i, \pi_{i+1})$  such that  $\pi_{i+1} = \pi_i - 1$ ;
- lrM**( $\pi$ ) = number of left-to-right maxima, i.e.  $i \geq 1$  such that  $\pi_i > \pi_j$  for all  $j < i$ ;
- rlm**( $\pi$ ) = number of right-to-left minima, i.e.  $i \geq 1$  such that  $\pi_i < \pi_j$  for all  $j > i$ ;
- inv**( $\pi$ ) = number of inversions, i.e. pairs  $(\pi_i, \pi_j)$  with  $\pi_i > \pi_j$  and  $i < j$ ,
- lmax**( $\pi$ ) = maximum value of the Lehmer code  $\ell_1 \ell_2 \dots \ell_n$  of  $\pi$ , i.e.  $\max_{1 \leq i \leq n} \ell_i$  where  $\ell_i = |\{\pi_j > \pi_i, j < i\}|$ ;
- lsum**( $\pi$ ) = sum of all values of the Lehmer code of  $\pi$ .

For a forest  $f \in \mathcal{F}_n$ , we define

- ledg**( $f$ ) = number of left edges, i.e., leftmost edges among its siblings;
- redg**( $f$ ) = number of right edges, i.e., rightmost edges among its siblings (**ledg**( $f$ ) = **redg**( $f$ ));
- nod1**( $f$ ) = number of nodes with only one child;
- ordt**( $f$ ) = number of ordered trees;
- leav**( $f$ ) = number of leaves, i.e., nodes without child;
- vpat**( $f$ ) = number of vertical paths (a vertical path is a path between a node and one of its ancestors);
- dept**( $f$ ) = depth, i.e., the maximal length of a vertical path;
- inpl**( $f$ ) = internal path length, i.e., the sum of the lengths of all paths from a node to the root.

In the following, the notation **st** will be refer to one of these statistics on the sets  $S_n(231)$  or  $\mathcal{F}_n$ . According to these definitions, it is straightforward to check that  $\phi$  transports these statistics as related in Table 2.

$S_n(231)$	<b>des = ides</b>	<b>adj</b>	<b>lrM</b>	<b>rlm</b>	<b>inv</b>	<b>lmax</b>	<b>lsum</b>
$\mathcal{F}_n$	<b>ledg = redg</b>	<b>nod1</b>	<b>ordt</b>	<b>leav</b>	<b>vpat</b>	<b>dept</b>	<b>inpl</b>

Table 2: Correspondences of statistics by the bijection  $\phi$  from  $S_n(231)$  to  $\mathcal{F}_n$ .

Using the correspondence between these statistics and taking advantage of the recursive structure of a forest, we derive several bivariate generating functions for two sets of pattern avoiding permutations with respect to the length and the statistics **st**.

**Theorem 4** For any statistic  $\mathbf{st}$  defined above, second column of Table 3 provides the bivariate generating function where the coefficient of  $z^n y^k$  is the number of permutations  $\pi \in S_n(231)$  with  $\mathbf{st}(\pi) = k$ .

*Proof.* Let  $F(x, y)$  be the generating function where the coefficient of  $z^n y^k$  is the number of forests  $f \in \mathcal{F}_n$  with  $\mathbf{st}(f) = k$ . Since a forest  $f \in \mathcal{F}_n$  is a collection of ordered trees, we have  $F(z, y) = \frac{1}{1-T(z, y)}$  where  $T(z, y)$  is the generating function for the number of ordered trees with respect to the length and the statistic  $\mathbf{st}$ . Now, using the fact that a nonempty ordered tree is a node connected to the roots of the trees of a forest, we easily derive functional equations for each statistic  $\mathbf{st}$ :

- for **ledg**, **redg** and **dept**,  $T(z, y) = zy(F(z, y) - 1) + z$ ;
- for **nod1**,  $T(z, y) = zyT(z, y) + z(F(z, y) - T(z, y))$ ;
- for **ordt**,  $T(z, y) = zyF(z, 1)$ ;
- for **leav**,  $T(z, y) = z(F(z, y) - 1) + zy$ ;
- for **vpat** and **inpl**,  $T(z, y) = z(F(zy, y) - 1) + z$ .

A simple calculation (using Maple for instance) completes the proof.  $\square$

**Theorem 5** The sets  $S_n(231, \underline{51\bar{4}23})$ ,  $n \geq 1$ , are enumerated by the number of single-source directed animals on the square lattice ([A005773](#), [18]), and for any statistic  $\mathbf{st}$ , third column of Table 3 provides the bivariate generating function where the coefficient of  $z^n y^k$  is the number of permutations  $\pi \in S_n(231, \underline{51\bar{4}23})$  with  $\mathbf{st}(\pi) = k$ .

*Proof.* Let  $\mathcal{G}_n \subset \mathcal{F}_n$  be the set of forests of ordered binary trees, *i.e.*, ordered trees where each node has at most two children. Let us prove that we have  $\phi^{-1}(\mathcal{G}_n) = S_n(231, \underline{51\bar{4}23})$ . Let  $\pi$  be a permutation in  $S_n(231, \underline{51\bar{4}23})$  and  $f_\pi = \phi(\pi)$ . Since  $\pi$  avoids  $\underline{51\bar{4}23}$ , any pattern  $\underline{4123}$  can be expanded into a pattern  $\underline{51423}$  which implies that the corresponding forest  $f_\pi$  does not contain a node with more than two children. Conversely, if the forest  $f_\pi$  belongs to  $\mathcal{G}_n$ , then the degree of any node of  $f$  is at most two. Let us suppose that  $\pi$  contains the pattern  $\underline{4123}$  on  $\pi_i \pi_{i+1} \pi_j \pi_k$ ,  $i + 1 < j < k$ . If there does not exist  $\ell$ ,  $i + 1 < \ell < j$  such that  $\pi_k < \pi_\ell < \pi_i$ , by construction the forest  $f_\pi$  has the node  $\pi_i$  with at least three children  $\pi_{i+1}$ ,  $\pi_j$  and  $\pi_k$ . So, any pattern  $\underline{4123}$  in  $\pi$  can be expanded into a pattern  $\underline{51423}$ , which proves that  $\pi$  avoids  $\underline{51\bar{4}23}$ .

Let  $G(x, y)$  be the generating function where the coefficient of  $z^n y^k$  is the number of forests  $f \in \mathcal{G}_n$  with  $\mathbf{st}(f) = k$ . Since a forest  $f \in \mathcal{G}_n$  is a collection of the ordered binary trees, we have  $G(z, y) = \frac{1}{1-R(z, y)}$  where  $R(z, y)$  is the generating function for the number of ordered binary trees with respect to the length and the parameter  $\mathbf{st}$ . Now, using the fact that a nonempty ordered binary tree is a node connected to the roots of at most two ordered binary trees, we can easily derive functional equations for each statistic  $\mathbf{st}$ :



- for **ledg**, **redg** and **dept**,  $R(z, y) = z + zyR(z, y) + zyR(z, y)^2$ ;
- for **nod1**,  $R(z, y) = z + zyR(z, y) + zR(z, y)^2$ ;
- for **ordt**,  $R(z, y) = zy + zyR(z, 1) + zyR(z, 1)^2$ ;
- for **leav**,  $R(z, y) = zy + zR(z, y) + zR(z, y)^2$ ;
- for **vpat** and **inpl**,  $R(z, y) = z + zR(z, y) + zR(z, y)^2$ .

A simple calculation (using Maple for instance) completes the proof.  $\square$

st	$S_n(231)$	$S_n(231, \underline{51\bar{4}23})$
{ }	Catalan ( <a href="#">A000108</a> , [18])	Directed animals ( <a href="#">A005773</a> , [18])
des, ides, lmax	$\frac{1-z+zy-\sqrt{z^2y^2-2z^2y+z^2-2zy-2z+1}}{2zy}$	$\frac{2yz}{3yz-1+\sqrt{y^2z^2-4yz^2-2yz+1}}$
adj	$\frac{1-zy+z-\sqrt{z^2y^2+2z^2y-3z^2-2zy-2z+1}}{2z}$	$\frac{2z}{2z-1+yz+\sqrt{y^2z^2-2yz-4z^2+1}}$
lrM	$\frac{2}{2-y+y\sqrt{1-4z}}$	$\frac{2z}{yz-y+2z+y\sqrt{-3z^2-2z+1}}$
rlm	$\frac{1+z-zy-\sqrt{z^2y^2-2z^2y+z^2-2zy-2z+1}}{2z}$	$\frac{2z}{3z-1+\sqrt{-4yz^2+z^2-2z+1}}$
inv, lsum	$F(z, y) = \frac{1}{1-z(F(zy, y)-1)-x}$	$\begin{cases} R(z, y) = z + zR(yz, y) + zR(yz, y)^2 \\ G(z, y) = \frac{1}{1-R(z, y)} \end{cases}$

Table 3: Generating functions for  $S_n(231)$  and  $S_n(231, \underline{51\bar{4}23})$  with respect to the length and the statistic **st**.

Notice that for **inv** and **lsum**, functional equations provide generating functions as continued fractions instead of closed forms.

## 4 Enumeration of $S_n(123)^\sim$

In this section we prove that the set  $S_n(123)^\sim$  is enumerated by the number of single-source directed animals on the square lattice ([A005773](#), [18]). To achieve this, we construct a bijection between  $S_n(123)^\sim$  and the set of forests of ordered binary trees, *i.e.*, trees where nodes have at most two ordered children (if a node has only one child then we call *0-edge* the corresponding link, and if a node has two children then the two corresponding links are called *0-edge* and *1-edge*, which defines an order on siblings).

A *run* of pure descents (also called *run* for short) in  $\pi = \pi_1 \dots \pi_n \in S_n$  is a maximal subsequence  $\pi_i \pi_{i+1} \dots \pi_j$ ,  $1 \leq i \leq j \leq n$ , of successive pure descents, *i.e.*  $(\pi_k, \pi_{k+1})$  is a pure descent for  $i \leq k \leq j-1$ , and the two pairs  $(\pi_{i-1}, \pi_i)$ ,  $(\pi_j, \pi_{j+1})$  are not pure descents (a run contains at least one entry, that is  $\pi_i$ ). To any run  $R$  of  $\pi \in S_n$ , we

associate the interval  $I(R) = [a, b] \subseteq [n]$  where  $a$  and  $b$  are the extremities of  $R$ , that is  $a = \min R$  and  $b = \max R$ .

In a permutation  $\pi \in S_n(123)$ , there do not exist three runs  $R, S$  and  $T$  such that  $I(R) \subset I(S) \subset I(T)$  (otherwise a pattern 123 would be created on the three entries  $\min T, \min S$  and  $\min R$ ). So, whenever there are two runs  $S, R$  such that  $I(S) \subset I(R)$ , we will say that  $S$  is a *secondary run*, and  $S$  appears necessarily in  $\pi$  at the right of  $R$ . A run  $R$  that is not secondary will be called *primary*. The family of intervals  $I(R)$  associated to the primary runs of  $\pi \in S_n(123)$  forms a partition of  $[n]$ . We denote by  $p \geq 1$  the cardinality of this partition, and let  $I_i, 1 \leq i \leq p$ , be the  $i$ th interval (considered in decreasing order), and let  $P_i$  be its associated primary run ( $I(P_i) = I_i$ ). For  $1 \leq i \leq p$ , let  $L_i$  be the restriction of  $\pi$  to the interval  $I_i$ . It can be decomposed as  $L_i = P_i S_i^1 S_i^2 \dots S_i^{s_i}$  where  $P_i$  is the  $i$ th primary run of  $\pi$ ,  $S_i^j$  is the  $j$ th secondary run of the interval  $I_i$  and  $s_i$  is the number of secondary runs in  $L_i$ .

We say there is a *break* between two consecutive secondary runs  $S_i^j$  and  $S_i^{j+1}$  in  $L_i$  if  $\min S_i^j = 1 + \max S_i^{j+1}$ . We refer to Figure 3 for an illustration of such a decomposition.




Figure 3: Illustration of the decomposition into runs of the permutation  $\pi = 12\ 8\ 14\ 13\ 7\ 5\ 3\ 11\ 10\ 2\ 1\ 9\ 6\ 4$ ;  $P_1 = 14\ 13$ ,  $P_2 = 12\ 8$ ,  $S_2^1 = 11\ 10$ ,  $S_2^2 = 9$ ,  $P_3 = 7\ 5\ 3$ ,  $S_3^1 = 6$ ,  $S_3^2 = 4$ ,  $P_4 = 2\ 1$ . A break occurs between the two consecutive secondary runs  $S_2^1$  and  $S_2^2$ .

With the above definitions we have Lemma 1.

**Lemma 1** *Let  $\pi \in S_n(123)$  and  $1 \leq i \leq p$ . If there is a break between two consecutive secondary runs  $S_i^j$  and  $S_i^{j+1}$  in  $L_i$ , then there exists a unique primary run  $P_k$  between  $S_i^j$  and  $S_i^{j+1}$  in the one-line notation of  $\pi$ , and we necessarily have  $k > i$ .*

*Proof.* If there is a break between  $S_i^j$  and  $S_i^{j+1}$ , then we have  $\min S_i^j = 1 + \max S_i^{j+1}$  and  $(\min S_i^j, \max S_i^{j+1})$  is not a pure descent. So, there exists an entry  $x$  of  $\pi$  between  $S_i^j$  and  $S_i^{j+1}$ , *i.e.*  $S_i^j$  and  $S_i^{j+1}$  are not contiguous. As  $\pi$  avoids 123,  $x$  is necessarily less than  $\min P_i$  where  $P_i$  is the primary run of  $L_i$ , and it does not belong to a secondary run. For a contradiction, let us assume that there is two entries  $x$  and  $y$ ,  $x > y$ , between  $S_i^j$  and  $S_i^{j+1}$ , that do not belong to the same primary run (we take  $y$  maximal such that  $x > y$ ). Obviously,  $x$  is on the left of  $y$ , otherwise it would create a pattern 123 on the entries  $yx \min S_i^{j+1}$ . Let  $P_k$  (resp.  $P_\ell$ ,  $\ell > k$ ) be the primary run that contains  $x$  (resp.  $y$ ). Since  $x > y$  and  $P_k$  is on the left of  $P_\ell$ , the two primary runs are not contiguous, and there exists a value  $z$  between  $P_k$  and  $P_\ell$ . The maximality of  $y$  implies that  $z$  is either below  $P_\ell$  or above  $P_k$ , which creates a pattern 123 in both cases. Thus, we obtain the desired contradiction.  $\square$

Lemma 1 allows us to define an injective map  $\alpha$  from the set  $\mathcal{B}$  of breaks to the set  $\mathcal{P}$  of primary runs, where the image by  $\alpha$  of a break is the unique primary run defined in Lemma 1. Moreover, it is easy to check that the map  $\alpha$  is increasing, *i.e.*, if  $B_1, B_2, \dots, B_r$  are the breaks of  $\mathcal{B}$  ordered in decreasing order (from top to bottom in the graphical representation of  $\pi$ ), then the two primary runs  $P_k = \alpha(B_i)$  and  $P_\ell = \alpha(B_j)$ ,  $1 \leq i < j \leq r$ , satisfy  $k < \ell$  (*i.e.*,  $P_k > P_\ell$ , which means that  $P_k$  is above  $P_\ell$  in the graphical representation of  $\pi$ ). The existence of this increasing map  $\alpha$  allows us to define another increasing map  $\beta$  (possibly equal to  $\alpha$ ) from  $\mathcal{B}$  to  $\mathcal{P}$ :

- $\beta(B_1)$  is the highest primary run below  $B_1$  (it always exists since  $\alpha(B_1)$  is below  $B_1$ );

- Let us assume that  $\beta$  is defined on  $U_i = \{B_1, \dots, B_i\}$ ,  $i \geq 1$ , and  $\beta$  is increasing such that  $\beta(B_j) \leq \alpha(B_j)$  for  $1 \leq j \leq i$ . Setting  $V_i = \beta(U_i)$ , we define  $\beta(B_{i+1})$  by the highest primary run below  $B_{i+1}$  that does not lie in  $V_i$  (it always exists since  $\alpha(B_{i+1})$  is below  $B_{i+1}$  and  $\alpha(B_{i+1})$  cannot lie in  $V_i$  since  $\alpha(B_{i+1})$  is below  $\alpha(B_i)$  and thus, also below  $\beta(B_i)$ ).

A crucial property of  $\beta$  is that it depends only on the set of primary and secondary runs, which means that two permutations in the same equivalence class provide the same map  $\beta$ .

Using the map  $\beta$ , we construct a forest  $\chi(\pi)$  of ordered binary trees from the graphical representation of  $\pi$  by adding 0-edges and 1-edges between some entries of  $\pi$  using the following process.

- (i) If  $(\pi_i, \pi_{i+1})$  is a pure descent, then we add a 0-edge between  $\pi_i$  and  $\pi_{i+1}$ .

- (ii) If there is break between  $S_i^j$  and  $S_i^{j+1}$ , then we add a 0-edge between  $\min S_i^j$  and  $\max S_i^{j+1}$ .
- (iii) If  $S_i^j$  is a secondary run and there is no break just before  $S_i^j$ , then we add a 1-edge between  $x$  and  $\max S_i^{j+1}$  where  $x$  is the smallest entry greater than  $\max S_i^{j+1}$  in the primary run  $P_i$ .
- (iv) If there is a break  $B$  between  $S_i^j$  and  $S_i^{j+1}$ , then we add a 1-edge between  $\max S_i^j$  and  $\max \beta(B)$ .

At the end of this process, we read the different connected components (rotated clockwise by  $\frac{\pi}{4}$ ) from top to bottom, and we draw the corresponding trees so that any 0-edge points to the left child and 1-edge points to the right child. See Figure 4 for an illustration of this construction. Black (resp. blue, red, green) edges come from (i) (resp. (ii), (iii), (iv)). In what follows, an edge  $e$  in  $\chi(\pi)$  will be denoted  $(a, b)$  where  $a$  and  $b$  are the extremities of  $e$  such that  $a$  is the parent of  $b$ .

Since any node in  $\chi(\pi)$  has at most two children,  $\chi(\pi)$  is a forest of ordered binary trees. Let  $T$  be a binary tree of  $\chi(\pi)$ . For any node  $v \in T$  we denote by  $r(v)$  (resp.  $l(v)$ ) the number of 1-edges (resp. 0-edges) in the path connecting the root of  $T$  with  $v$ . We say that a node  $v$  is isolated when it has no siblings.

**Remark 1** *Let  $e = (a, b)$  be a 0-edge in  $\chi(\pi)$ .*

- $(a, b)$  is a pure descent in a primary run of  $\pi$  if and only if  $r(a)$  is even.
- $(a, b)$  is a pure descent in a secondary run of  $\pi$  if and only if  $a$  is isolated and  $r(a)$  is odd.
- There is a break between  $a$  and  $b$  if and only if  $a$  is not isolated and  $r(a)$  is odd.

Denoting by  $\text{rath}(v)$  the binary word consisting of edge labels in the path from the root to  $v$  and using a lexicographical order over such binary words (e.g.  $101 > 011$ ,  $1 > 01$ ), we define a total order on the set of nodes  $V$  in  $\chi(\pi)$ . For  $a, b \in V$ , we set

$$a < b \iff \begin{cases} \text{either } a \text{ belongs to a tree before that of } b \text{ in the forest } \chi(\pi), \\ \text{or } \begin{cases} r(a) < r(b) \text{ or} \\ r(a) = r(b) \text{ and } \text{rath}(a) > \text{rath}(b) \end{cases} \end{cases} \quad (\star)$$

where  $r(a)$  and is defined before Remark 1. We extend this order for paths  $v_1v_2\dots$  where  $v_i$  and  $v_{i+1}$  are nodes of  $\chi(\pi)$  connected by a 0-edge: two disjoint paths  $v_1v_2v_3\dots$  and  $u_1u_2u_3\dots$  are compared by their heads, i.e.

$$(v_1v_2v_3\dots) < (u_1u_2u_3\dots) \iff v_1 < u_1.$$

Extracting from the forest  $\chi(\pi)$  certain subsets of disjoint paths and taking into account the above order relation, we obtain the following.

**Remark 2** *The three statements hold:*

- *The  $i$ th primary run  $P_i$  in  $\pi$  (ordered from the top) corresponds to the  $i$ th maximal path  $v_1v_2v_3\dots$  of consecutive nodes joined by 0-edges in  $\chi(\pi)$  where  $r(v_1)$  is even.*
- *The  $i$ th secondary run in  $\pi$  (ordered from the top) corresponds to the  $i$ th maximal path  $v_1v_2v_3\dots$  of consecutive isolated nodes joined by 0-edges in  $\chi(\pi)$  where  $r(v_1)$  is odd.*
- *The  $i$ th break in  $\pi$  (ordered from the top) corresponds to the  $i$ th 0-edge  $(a, b)$  in  $\chi(\pi)$  such that  $r(a)$  is odd and  $a$  is isolated.*

**Consequence:** Let  $(a, b)$  be a pure descent in a primary run of  $\pi$ , and  $e = (a, b)$  its associated 0-edge in  $\chi(\pi)$ . Then, the number of entries of  $\pi$  in the interval  $]b, a[$ , *i.e.*  $a - b - 1$ , is equal to the number of nodes in the maximal path of 0-edges starting on the right child of  $a$ .




Figure 4: The permutation  $\pi = 12\ 8\ 14\ 13\ 7\ 5\ 3\ 11\ 10\ 2\ 1\ 9\ 6\ 4$  and the corresponding forest with nodes labeled using  $(\star)$  order relation.

**Proposition 1** *Let  $\pi$  and  $\pi'$  be two permutations in  $S_n(123)$ .*

- *If  $\pi$  and  $\pi'$  belong to the same equivalence class, then  $\chi(\pi) = \chi(\sigma)$ .*
- *If  $\pi$  and  $\pi'$  belong to different equivalence classes, then  $\chi(\pi) \neq \chi(\sigma)$ .*

*Proof.* As we have seen above, the map  $\beta$  depends only on the set of primary and secondary runs. Thus, our construction applied on two permutations lying in the same class provides the same forest.

Moreover, if two permutations  $\pi$  and  $\pi'$  do not belong to the same class then their sets of primary and secondary runs necessarily differ. Due to the statements of Remark 2 and the above consequence, we deduce easily that  $\chi(\pi)$  and  $\chi(\pi')$  are different.  $\square$

**Theorem 6** *The sets  $S_n(123)^\sim$ ,  $n \geq 1$ , are enumerated by the numbers of single-source directed animals on the square lattice (A005773, [18]).*

*Proof.* Proposition 1 proves that  $\chi$  is injective. So, it suffices to show the surjectivity of  $\chi$ , *i.e.*, any forest of ordered binary trees is the image by  $\chi$  of a permutation avoiding 123.

First, we prove that any binary tree can be obtained from a permutation  $\pi \in S_n(123)$  by the above construction. Let  $T$  be a binary tree with  $n$  nodes. By Remark 2, a maximal path  $P$  of nodes connected by 0-edges in  $T$  such that  $r(P)$  is even corresponds to a primary run of  $\pi$ . Moreover, if a 0-edge  $e$  corresponds to a pure descent  $(a, b)$  in a primary run of  $\pi$ ,  $b(e) = a - b - 1$  is the number of nodes in the maximal path (possibly reduced to one node) of 0-edges starting on the right child of  $a$ .

Then the primary runs of  $\pi$  are entirely determined by the sequence  $b_1 b_2 \dots b_k$  with  $b_1 = n$  and  $b_i = b(e)$  where  $e$  is the  $i$ th 0-edge of  $T$  (using the  $(\star)$  order relation) such that  $r(e)$  is even. If  $p$  is the number of primary runs and  $P_i$  is the  $i$ th primary run of  $\pi$  then the sequence  $P_1 P_2, \dots, P_p$  is decreasing.

Consequently, secondary runs of  $\pi$  take values from  $[n] \setminus \cup_i^p P_i$ , and the breaks correspond to the 0-edges  $e = (a, b)$  where  $a$  is non-isolated and  $r(a)$  is odd, which entirely determines secondary runs and breaks. If  $q$  is the number of secondary runs and  $S_i$  is the  $i$ th secondary run (using the  $(\star)$  order relation) then the sequence  $S_1 S_2 \dots S_q$  is decreasing.

Now we construct a permutation  $\pi$  avoiding 123 by a shuffle of the two decreasing sequences  $P_1 P_2, \dots, P_p$  and  $S_1 S_2 \dots S_q$ . We read  $P_1 S_1 S_2 \dots S_q$  from left to right, and whenever we meet a break between  $S_i$  and  $S_{i+1}$  we insert between them the first primary run not yet inserted (this is exactly the correspondence given by the increasing map  $\beta$  defined above). Obviously, the sequence obtained at the end of the process is a permutation avoiding the pattern 123 since it is a shuffle of two decreasing sequences.

Finally, the image of  $\pi$  by  $\chi$  provides the tree  $T$ , which means that  $\chi(S_n(123))$  contains the set of all ordered binary trees of size  $n$ .

So, let us assume that  $f$  is a forest of ordered binary trees  $T_1, T_2, \dots, T_k$ . For  $1 \leq i \leq k$ , we construct the permutation  $\pi_i$  from the tree  $T_i$  by the previous process, *i.e.*,  $\pi_i = \chi^{-1}(T_i)$ . Let  $\pi$  be the permutation obtained by the skew sum  $\pi_1 \ominus \pi_2 \ominus \dots \ominus \pi_k$  where  $\pi \ominus \pi'$  is the permutation  $\sigma$  such that

$$\sigma(i) = \begin{cases} \pi(i) + m' & \text{for } 1 \leq i \leq m, \\ \pi'(i - m) & \text{for } m + 1 \leq i \leq m + m' \end{cases}$$

where  $m$  (resp.  $m'$ ) is the length of  $\pi$  (resp.  $\pi'$ ).

Now we read  $\pi_1 \ominus \pi_2 \ominus \dots \ominus \pi_k$  from left to right. Whenever a non desirable pure descent have been created between  $\pi_i$  and  $\pi_{i+1}$ , it is easy to see that  $\pi_i$  is necessarily a decreasing sequence. In this case we permute  $\pi_i$  and the first primary run of  $\pi_{i+1}$ . At the end of the process, the permutation  $\pi$  satisfies  $\chi(\pi) = f$ .  $\square$

For instance, the previous construction applied on the forest illustrated in Figure 4 provides the permutation  $\pi = 12\ 8\ 14\ 13\ 11\ 10\ 7\ 5\ 3\ 9\ 6\ 4\ 2\ 1$ . Indeed, we have  $\pi_1 = \chi^{-1}(T_1) = 2\ 1$ ,  $\pi_2 = \chi^{-1}(T_2) = 10\ 6\ 9\ 8\ 5\ 3\ 1\ 7\ 4\ 2$  and  $\pi_3 = \chi^{-1}(T_3) = 2\ 1$ . Since a non desirable pure descent  $(13, 12)$  is created in the permutation  $\pi_1 \ominus \pi_2 \ominus \pi_3 = \mathbf{14\ 13}\ 12\ 8\ 11\ 10\ 7\ 5\ 3\ 9\ 6\ 4\ \mathbf{2\ 1}$ , we permute the two primary runs  $12\ 8$  and  $14\ 13$ , which gives  $\pi = 12\ 8\ \mathbf{14\ 13}\ 11\ 10\ 7\ 5\ 3\ 9\ 6\ 4\ 2\ 1$ .

## 5 Going further

We conclude this paper by giving several open questions and possible research directions.

We obtain experimentally the numbers of classes in  $S_n(132)^\sim$  and  $S_n(213)^\sim$  for small values of  $n$ ,  $1 \leq n \leq 9$ . For  $S_n(132)^\sim$ , we obtain the sequence  $1, 2, 4, 10, 26, 66, 169, 437, 1130$  and for  $S_n(213)^\sim$ , we obtain the sequence  $1, 2, 4, 9, 22, 56, 146, 388, 1048$ . The first sequence does not appear in [18], while the second sequence seems to be  $A152225$  which corresponds to the number of Dyck paths of semilength  $n$  with no peaks at height  $0 \pmod 3$  and no valleys at height  $2 \pmod 3$ . Is it possible to obtain the generating functions for these sets and to make a link with Dyck paths?

In [2], the authors give a one-to-one correspondence between the set  $\mathcal{F}_n$  of forests of ordered trees and the set  $S_n(321, 4\bar{1}523)$  that transports various parameters. However, they do not succeed to give an interpretation for the number of inversions, the degree of the root less one and the internal-path-length. In Section 2, we exhibit a bijection between  $\mathcal{F}_n$  and  $S_n(231, \underline{51}\bar{4}23)$ , which gives a new set of pattern avoiding permutations

enumerated as the single-source directed animals on the square lattice. This bijection has the advantage that it transports many parameters (see Section 3), and in particular the three previous parameters. Is it possible to give an interpretation of these parameters in term of the single-source directed animals?

In Section 4, we prove that  $S_n(123)^\sim$  is enumerated by the number of directed animals (or equivalently directed polyominoes). Is it possible to give an interpretation of the equivalence relation in term of polyominoes?

## References

- [1] E. Babson and E. Steingrímsson. Generalized permutation patterns and a classification of Mahonian statistics. *Sém. Lothar. Combin.* (electronic), 44, 2000.
- [2] E. Barucci, A. Del Lungo, E. Pergola and R. Pinzani. Directed animals, forests and permutations. *Discrete Mathematics*, 204, 41–71, 1999.
- [3] J.-L. Baril, T. Mansour and A. Petrossian. Equivalence classes of permutations modulo excedances. *Journal of Combinatorics*, 5(4), 453–469, 2014.
- [4] J.-L. Baril and A. Petrossian. Equivalence classes of permutations modulo descents and left-to-right maxima. *Pure Mathematics and Applications*, 25(1), 19–29, 2015.
- [5] J.-L. Baril and S. Kirgizov. The pure descent statistics in permutations. *To be published in Discrete Mathematics*, 2016.
- [6] F. Bergeron, N. Bergeron, R.B. Howlett and D.E. Taylor. A decomposition of the descent algebra of a finite Coxeter group. *J. Algebraic Combin.*, 1, 23–44, 1992.
- [7] M. Bóna. *Combinatorics of permutations*. Chapman and Hall/CRC, Second Edition, 2012.
- [8] M. Bousquet-Mélou. New enumerative results on two-dimensional directed animals. *Proc. 7th FPSAC*, Université de Marne-la-Vallée, Paris, 1995, 103–116.
- [9] I. Gessel and G. Viennot. Binomial determinants, paths, and hook length formulae. *Adv. Math.*, 58, 300–321, 1985.
- [10] S. Kitaev. *Patterns in permutations and words*. Springer-Verlag, 2011.
- [11] D.E. Knuth. *The art of computer programming*. Vol. 1, Addison-Wesley, Reading MA, 1973, Third edition, 1997.



- [12] P.A. MacMahon. *Combinatory Analysis*. Vols. 1 and 2, Cambridge Univ. Press, Cambridge, UK, 1915 (reprinted by Chelsea, New-York, 1955).
- [13] T. Mansour. *Combinatorics of Set Partitions*. *Discrete Mathematics and its Applications*, Chapman & Hall/CRC, 2013.
- [14] T. Mansour. Permutations avoiding a pattern from  $S_k$  and at least two patterns from  $S_3$ . *Ars. Combin.*, 62, 227–239, 2002.
- [15] L. Pudwell. Enumeration schemes for permutations avoiding barred patterns. *The Electronic Journal of Combinatorics*, 17, R29, 2010.
- [16] T. Mansour and A. Vainshtein. Restricted 132-avoiding permutations. *Adv. Appl. Math.*, 26, 258–269, 2001.
- [17] R. Simion and F.W. Schmidt. Restricted permutations. *European Journal of Combinatorics*, 6, 383-406, 1985.
- [18] N.J.A. Sloane. On-Line Encyclopedia of Integer Sequences. Published electronically at <http://oeis.org/>.
- [19] R. Stanley. *Enumerative Combinatorics*. *Cambridge University Press*, Vol. 1, 1997.
- [20] J. West. Permutations with forbidden subsequences and stack-sortable permutations. Ph.D. Thesis, MIT, 1990.
- [21] J. West. Generating trees and forbidden subsequences. *Proceeding of sixth FPSAC*, 441–450, 1994.
- [22] J. West. Generating trees and the Catalan and Schröder numbers. *Discrete Math.*, 146, 247–262, 1995.
- [23] R.M. Wilson and J.H. Lint. *A course in combinatorics*. Vol. I, Cambridge University Press, 2002.