

Transformation à la Foata for special kinds of descents and excedances

Jean-Luc Baril and Sergey Kirgizov

LIB, Université Bourgogne Franche-Comté
B.P. 47 870, 21078 Dijon-Cedex, France
[barjl](mailto:barjl@u-bourgogne.fr), [sergey.kirgizov](mailto:sergey.kirgizov@u-bourgogne.fr)@u-bourgogne.fr

January 6, 2021

Abstract

A pure excedance in a permutation $\pi = \pi_1\pi_2\dots\pi_n$ is a position $i < \pi_i$, $1 \leq i \leq n-1$, so that there is no $j < i$ such that $i \leq \pi_j < \pi_i$. We present a one-to-one correspondence on the symmetric group that transports pure excedances to descents of special kind. As a byproduct, we prove that the popularity of pure excedances equals those of pure descents on permutations, while their distributions are different.

Keywords: Permutation, statistic, distribution, popularity, descent, excedance, cycle.

1 Introduction and notations

The distribution of the number of descents has been widely studied on several classes of combinatorial objects such as permutations [14], cycles [7, 8], and words [3, 10]. Many interpretations of this statistic appear in several fields as Coxeter groups [4, 11] or lattice theory [5, 12]. One of the most famous result involves the *Foata fundamental transformation* [9] to establish a one-to-one correspondence between descents and excedances on permutations. This bijection provides a more straightforward proof than those of MacMahon [14] for the equidistribution of these two Eulerian statistics.

In this paper, we present a bijection *à la Foata* on the symmetric group that exchanges pure excedances with special kind of descents defined as a mesh pattern p_2 [6] (see below for the definitions of these patterns). Then,

we deduce that the popularities but not the distributions of pure descents [2] and pure excedances are the same. They are given by the generalized Stirling number $n! \cdot (H_n - 1)$ (see A001705 in [15]) where $H_n = \sum_{k=1}^n \frac{1}{k}$ is the n -th harmonic number. Finally, we conjecture the existence of a bijection on the symmetric group that exchanges pure excedances and p_2 while preserving the number of cycles.

Let S_n be the set of permutations of length n , *i.e.*, all bijections from $[n] = \{1, 2, \dots, n\}$ into itself. The one-line representation of a permutation $\pi \in S_n$ is $\pi = \pi_1 \pi_2 \dots \pi_n$ where $\pi_i = \pi(i)$, $1 \leq i \leq n$. For $\sigma \in S_n$, the *product* $\sigma \cdot \pi$ is the permutation $\sigma(\pi_1) \sigma(\pi_2) \dots \sigma(\pi_n)$. A ℓ -*cycle* $\pi = \langle i_1, i_2, \dots, i_\ell \rangle$ is a n -length permutation satisfying $\pi(i_1) = i_2, \pi(i_2) = i_3, \dots, \pi(i_{\ell-1}) = i_\ell, \pi(i_\ell) = i_1$ and $\pi(j) = j$ for $j \in [n] \setminus \{i_1, i_2, \dots, i_\ell\}$. For $1 \leq k \leq n$, we denote by $C_{n,k}$ the set of all n -length permutations admitting a decomposition in a product of k disjoint cycles. The set $C_{n,k}$ is counted by the signless Stirling numbers of the first kind $c(n, k)$ defined by

$$c(n, k) = (n - 1) c(n - 1, k) + c(n - 1, k - 1)$$

where $c(n, k) = 0$ if $n \leq 0$ or $k \leq 0$, except $c(0, 0) = 1$ (see [16, 17]). These numbers also enumerate n -length permutations π having k *left-to-right maxima*, *i.e.*, values $i \in [n]$ such that $\pi_j < \pi_i$ for $j < i$ (see [16]), and permutations $\pi \in S_n$ with $k - 1$ *pure descents*, *i.e.*, descents $\pi_i > \pi_{i+1}$ where there is no $j < i$ such that $\pi_j \in [\pi_{i+1}, \pi_i]$ (see [2]). Note that a pure descent can be viewed as an occurrence of the mesh pattern $(21, L_1)$ where $L_1 = \{1\} \times [0, 2] \cup \{(0, 1)\}$. Indeed, for a k -length permutation σ and a subset $R \subseteq [0, k] \times [0, k]$, an occurrence of the mesh pattern (σ, R) in a permutation π is an occurrence of σ in π with the additional restriction that no element of π lies inside the shaded regions defined by R , where $(i, j) \in R$ means the square having bottom left corner (i, j) in the graphical representation $\{(i, \sigma_i), i \in [k]\}$ of σ . For instance, an occurrence of the mesh pattern p_1 in Figure 1 corresponds to an occurrence of a pure descent. See [6] for a more detailed definition of mesh patterns.

Regarding this interpretation of pure descents in terms of mesh patterns, we define other kinds of descents by the mesh patterns $p_i = (21, L_i)$, $p'_i = (21, R_i)$ with $L_i = \{1\} \times [0, 2] \cup \{(0, i)\}$ and $R_i = \{1\} \times [0, 2] \cup \{(2, i)\}$ for $0 \leq i \leq 2$. Modulo the fundamental symmetries on permutations (reverse and complement), it is straightforward to see that p_0, p_1 , and p_2 are respectively in the same distribution class as p'_2, p'_1 and p'_0 . Then, we deal with only mesh patterns $p_i, i \in [0, 2]$. We refer to Figure 1 for a graphical illustration. On the other hand, we define a *pure excedance* as an occurrence of an excedance,

i.e. $\pi_i > i$, with the additional restriction that there is no point (j, π_j) such that $1 \leq j \leq i - 1$ with $i \leq \pi_j < \pi_i$. Although such a pattern (called pe_x) is not a mesh pattern, we can represent it graphically as shown in Figure 1.




Figure 1: Illustration of the mesh patterns p_0 , p_1 , p_2 and pe_x ; p_1 and pe_x correspond respectively to a pure descent and a pure excedance.

A *statistic* is an integer-valued function from a set \mathcal{A} of n -length permutations (we use the boldface to denote statistics). For a pattern p , we define the pattern statistic $\mathbf{p} : \mathcal{A} \rightarrow \mathbb{N}$ where the image $\mathbf{p} \pi$ of $\pi \in \mathcal{A}$ by \mathbf{p} is the number of occurrences of p in π . The *popularity* of p in \mathcal{A} is the total number of occurrences of p over all objects of \mathcal{A} , that is $\sum_{a \in \mathcal{A}} \mathbf{p} a$ (see [5] for instance). Below, we present statistics that we use throughout the paper:

- exc** π = number of excedances in π ,
- pex** π = number of pure excedances in π ,
- des** π = number of descents in π ,
- des _{i}** π = number of patterns p_i in π , $0 \leq i \leq 2$,
- fix** π = number of fixed points in π ,
- cyc** π = number of cycles in the decomposition of π ,
- pcyc** π = number of pure cycles (i.e. cycles of length at least two) in π ,
= **cyc** π - **fix** π

We organize the paper as follows. In Section 2, we focus on patterns p_i , $0 \leq i \leq 2$. We prove that the statistics **des**₀ and **des**₁ are equidistributed by giving algebraic and bijective proofs. Next, we provide the bivariate exponential generating function for the distribution of p_2 , and we deduce that p_2 has the same popularity as p_0 and p_1 , without having the same distribution. In Section 3, we present a bijection on S_n that transports pure excedances into patterns p_2 . Notice that the Foata's first transformation is not a candidate for such a bijection. As a consequence, pure descents and pure excedances are equipopular on S_n , but they do not have the same distribution. Combining all these results, we deduce that patterns p_i , $0 \leq i \leq 2$, and pe_x are equipopular on the symmetric group S_n . Finally we present two conjectures about the

equidistribution of $(\mathbf{cyc}, \mathbf{des}_2)$ and $(\mathbf{cyc}, \mathbf{pex})$, and that of $(\mathbf{des}, \mathbf{des}_2)$ and $(\mathbf{exc}, \mathbf{pex})$.

2 The statistics \mathbf{des}_i , $0 \leq i \leq 2$

For $0 \leq i \leq 2$, let $A_{n,k}^i$ be the set of n -length permutations having k occurrences of p_i , and denote by $a_{n,k}^i$ its cardinality. Let $A^i(x, y)$ be the bivariate exponential generating function $\sum_{n=0}^{\infty} \sum_{k=0}^{n-1} a_{n,k}^i \frac{x^n}{n!} y^k$. In [2, 13], it is proved that $a_{n,k}^1$ equals the signless Stirling numbers of the first kind $c(n, k+1)$ (see A132393 in [15]). Indeed, a permutation $\sigma \in A_{n,k}^1$ can be uniquely obtained from an $(n-1)$ -length permutation π by one of the two following constructions:

- (i) if $\pi \in A_{n-1,k-1}^1$, then we increase by one all values of π greater than or equal to π_{n-1} , and we add π_{n-1} at the end;
- (ii) if $\pi \in A_{n-1,k}^1$, then we increase by one all values of π greater than or equal to a given value $x \leq n$, $x \neq \pi_{n-1}$ and we add x at the end.

Then, we deduce the recurrence relation $a_{n,k}^1 = a_{n-1,k-1}^1 + (n-1)a_{n-1,k}^1$ with $a_{n,0}^1 = (n-1)!$ for $n \geq 1$, $a_{0,0}^1 = 1$ and the bivariate exponential generating function is

$$A^1(x, y) = \frac{1}{y(1-x)^y} - \frac{1}{y} + 1$$

which proves that $a_{n,k}^1 = c(n, k+1)$.

Below, we prove that $a_{n,k}^1$ also counts n -length permutations having k occurrences of the pattern p_0 .

Theorem 1. *The number $a_{n,k}^0$ of n -length permutations having k occurrences of pattern p_0 equals $a_{n,k}^1 = c(n, k+1)$.*

Proof. An n -length permutation $\sigma \in A_{n,k}^0$ can be uniquely obtained from an $(n-1)$ -length permutation π by one of the two following constructions:

- (i) if $\pi \in A_{n-1,k-1}^0$, then we increase by one all values of π and we add 1 at the end;
- (ii) if $\pi \in A_{n-1,k}^0$, then we increase by one all values of π greater than or equal to given value x , $1 < x \leq n$, and we add x at the end.

We deduce the recurrence relation $a_{n,k}^0 = a_{n-1,k-1}^0 + (n-1)a_{n-1,k}^0$ with the initial condition $a_{n,0}^0 = (n-1)!$, and then $a_{n,k}^0 = a_{n,k}^1 = c(n, k+1)$. \square

Now, we focus on the distribution of the pattern p_2 . Table 1 provides exact values for small sizes.

Theorem 2. *We have*

$$A^2(x, y) = \frac{e^{x(1-y)}}{(1-x)^y},$$

and the general term $a_{n,k}^2$ satisfies for $n \geq 2$ and $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$

$$a_{n,k}^2 = na_{n-1,k}^2 + (n-1)a_{n-2,k-1}^2 - (n-1)a_{n-2,k}^2$$

with the initial conditions $a_{n,0}^2 = 1$ and $a_{n,k}^2 = 0$ for $n \geq 0$ and $k > \lfloor \frac{n}{2} \rfloor$ (see Table 1 and the triangular table A136394 in [15]).

Proof. Let $\sigma = \sigma_1\sigma_2 \dots \sigma_n$ denote a permutation of length n having k occurrences of pattern p_2 . Let $u_{n,k}$ (resp. $v_{n,k}$) be the number of such permutations satisfying $\sigma_n = n$ (resp. $\sigma_n < n$). Obviously, we have

$$a_{n,k}^2 = u_{n,k} + v_{n,k}.$$

A permutation σ with $\sigma_n = n$ can be uniquely constructed from an $(n-1)$ -length permutation π as $\sigma = \pi_1\pi_2 \dots \pi_{n-1}n$. No new occurrences of p_2 are created, and we obtain

$$u_{n,k} = a_{n-1,k}^2.$$

A permutation σ satisfying $\sigma_n < n$ can be uniquely obtained from an $(n-1)$ -length permutation π by adding a value $x < n$ on the right side of its one-line notation, after increasing by one all the values greater than or equal to x . This construction creates a new pattern p_2 if and only if π ends with $n-1$. Thus, we deduce

$$v_{n,k} = (n-1)u_{n-1,k-1} + (n-1)v_{n-1,k}.$$

Combining the equations, we obtain for $n \geq 2$ and $k \geq 1$

$$a_{n,k}^2 = na_{n-1,k}^2 + (n-1)a_{n-2,k-1}^2 - (n-1)a_{n-2,k}^2,$$

which implies the following differential equation

$$\frac{\partial A^2(x, y)}{\partial x} = (y-1)xA^2(x, y) + \frac{\partial (xA^2(x, y))}{\partial x}, \text{ where } A^2(x, 0) = 1.$$

A simple calculation provides the claimed closed form for the generating function $A^2(x, y)$. \square

Corollary 1. For $0 \leq i \leq 2$, the patterns p_i are equipopular on S_n . Their popularity is given by the generalized Stirling number $n! \cdot (H_n - 1)$ (see [A001705](#) in [15]) where $H_n = \sum_{k=1}^n \frac{1}{k}$ is the n -th harmonic number.

Proof. The generating function of the popularity is directly deduced from the bivariate generating function of pattern distribution by calculating

$$\left. \frac{\partial A^1(x, y)}{\partial y} \right|_{y=1} = \left. \frac{\partial A^2(x, y)}{\partial y} \right|_{y=1}.$$

□

| $k \setminus n$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|-----------------|---|---|---|----|-----|-----|------|-------|
| 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 1 | | 1 | 5 | 20 | 84 | 409 | 2365 | 16064 |
| 2 | | | | 3 | 35 | 295 | 2359 | 19670 |
| 3 | | | | | | 15 | 315 | 4480 |
| 4 | | | | | | | | 105 |
| ... | | | | | | | | ... |
| Σ | 1 | 2 | 6 | 24 | 120 | 720 | 5040 | 40320 |

Table 1: Number of n -length permutations having k occurrences of p_2 for $0 \leq k \leq 4$ and $1 \leq n \leq 8$.

The statistic \mathbf{des}_2 has a different distribution from \mathbf{des}_0 and \mathbf{des}_1 , but the three patterns p_0, p_1, p_2 have the same popularity. Below we present a bijection on S_n that transports the statistic \mathbf{des}_2 to the statistics $\mathbf{pcyc} = \mathbf{cyc} - \mathbf{fix}$.

Theorem 3. There is a one-to-one correspondence ϕ on S_n such that for any $\pi \in S_n$, we have

$$\mathbf{des}_2 \pi = \mathbf{pcyc} \phi(\pi).$$

Proof. Let π be a permutation of length n having k occurrences of p_2 . We decompose

$$\pi = B_0 \pi_{i_1} A_1 B_1 \pi_{i_2} A_2 B_2 \pi_{i_3} \dots \pi_{i_k} A_k B_k,$$

where

- $\pi_{i_1} < \pi_{i_2} < \dots < \pi_{i_k}$ are the tops of the occurrences of p_2 , *i.e.* values $\pi_{i_j} > \pi_{i_{j+1}}$ such that there does not exist $\ell < i_j$ such that $\pi_\ell > \pi_{i_j}$,
- A_j is a maximal sequence such that all its values are lower than π_{i_j} ,
- for $0 \leq j \leq k$, B_j is an increasing sequence such that $\pi_{i_j} < \min B_j$ and $\max B_j < \pi_{i_{j+1}}$.

Now we construct an n -length permutation $\phi(\pi)$ with k pure cycles as follows:

$$\phi(\pi) = \langle \pi_{i_1} A_1 \rangle \cdot \langle \pi_{i_2} A_2 \rangle \cdots \langle \pi_{i_k} A_k \rangle.$$

For instance, if $\pi = 125346879$ then $\phi(\pi) = \langle 5, 3, 4 \rangle \cdot \langle 8, 7 \rangle$. The map ϕ is clearly a bijection on S_n such that $\mathbf{des}_2 \pi$ equals the number of pure cycles in $\phi(\pi)$. \square

Note that ϕ^{-1} is closely related to the Foata fundamental transformation.

3 The statistic \mathbf{pex} of pure excedances

In order to prove the equidistribution of \mathbf{pex} and \mathbf{des}_2 , regarding Theorem 3, it suffices to construct a bijection on S_n that transports pure excedances to pure cycles. Here, we first exhibit a bijection on the set D_n of n -length derangements (permutations without fixed points), then we extend it to the set of all permutations S_n .

Any permutation $\pi \in S_n$ is uniquely decomposed as a product of transpositions of the following form:

$$\pi = \langle t_1, 1 \rangle \cdot \langle t_2, 2 \rangle \cdots \langle t_n, n \rangle$$

where t_i are integers such that $1 \leq t_i \leq i$. The transposition array of π is defined by $T(\pi) = t_1 t_2 \dots t_n$, which induces a bijection $\pi \mapsto T(\pi)$ from S_n to the product set $T_n = [1] \times [2] \times \dots \times [n]$. By Lemma 1 from [1], the number of cycles of a permutation π is given by the number of fixed points in $T(\pi)$. Moreover, it is straightforward to check the two following properties:

- if $t_i = i$, then $\pi_i = i$ if and only if there is no number $j > i$ such that $t_j = t_i = i$;
- if $t_i = i$ and $\pi_i \neq i$, then i is the minimal element of a cycle of length at least two in π .

So, we deduce the following lemma.

Lemma 1. *The transposition array $t_1 t_2 \dots t_n \in T_n$ corresponds to a derangement if and only if: $t_i = i \Rightarrow$ there is $j > i$ such that $t_j = i$.*

Given a derangement $\pi = \pi_1\pi_2 \dots \pi_n \in D_n$ and its graphical representation $\{(i, \pi_i), i \in [n]\}$. We say that the square $(i, j) \in [n] \times [n]$ is *free* if all following conditions hold:

- (i) Neither π_i nor i is a position of a pure excedance;
- (ii) (i, j) is not on the first diagonal, *i.e.* $j \neq i$;
- (iii) there does not exist $k > i$ such that $\pi_k = j$;
- (iv) j is not a pure excedance such that $j < i$ and $\pi^{-1}(j) < i$;
- (v) there does not exist $k < i$, with $\pi_k = j > i$ such that all values of the interval $[i, j - 1]$ appear on the right of π_i in π .

Whenever at least one of the statements above is not satisfied, we say that the square (i, j) is *unfree*. Notice that if i and π_i are not the positions of a pure excedance, then the square (i, π_i) is always free. So, for a column i of the graphical representation of π such that i and π_i are not the positions of a pure excedance, we label by j the j th free square from the bottom to the top. We refer to Figure 2 for an example of this labelling.

Now we define the map λ from D_n to the set T_n^\bullet of transposition arrays of length n satisfying the property of Lemma 1.

For a permutation $\pi = \pi_1\pi_2 \dots \pi_n \in D_n$, we label its graphical representation as defined above, and $\lambda(\pi) = \lambda_1\lambda_2 \dots \lambda_n$ is obtained as follows:

- if i is a pure excedance in π , then we set $\lambda_i = i$ and $\lambda_{\pi^{-1}(i)} = i$;
- otherwise, λ_i is the sum of the label of the free square (i, π_i) with the number of pure excedances $k < i$ such that $\pi^{-1}(k) < i$.

For instance, if $\pi = 6\ 8\ 12\ 5\ 4\ 7\ 3\ 2\ 11\ 1\ 9\ 10$ then we obtain $\lambda(\pi) = 1\ 1\ 2\ 4\ 4\ 2\ 1\ 1\ 9\ 1\ 9\ 10$ (see Figure 2).

Let us consider i , $1 \leq i \leq n$. If i is a pure excedance of π , then we fix $\lambda_i = i$ and $\lambda_{\pi^{-1}(i)} = i < \pi^{-1}(i)$. Otherwise, the square (i, i) is unfree, and all squares (i, π_k) , $i + 1 \leq k \leq n$, are unfree, which implies that the number of free squares in the i th column is less than or equal to i . This means that $\lambda(\pi)$ lies in T_n . Note that, by construction, all labeled squares do not correspond to any pure excedance. Now let us prove that the square (i, π_i) cannot be labeled i . Indeed, if $\pi_i < i$ then the label of (i, π_i) is necessarily at most $\pi_i \leq i - 1$; otherwise, if $\pi_i > i$ then the fact that i is not a pure excedance implies that there is $\pi_j \in [i, \pi_i - 1]$ with $j < i$. Let us choose the lowest j with this property. Using (v), the square (i, j) is unfree, which implies that

the label of (i, π_i) is less than or equal to n minus the minimal number of unfree squares (i, j) in column i , that is $n - (n - i + 1) = i - 1$. Moreover, the transposition array $\lambda(\pi)$ has exactly **pex** π fixed points, and for any fixed point i there necessarily exists $j = \pi^{-1}(i) > i$ such that $\lambda_j = \lambda_i = i$. This implies that $\lambda(\pi) \in T_n^\bullet$.




Figure 2: Illustration of the bijection λ for $\pi = 6\ 8\ 12\ 5\ 4\ 7\ 3\ 2\ 11\ 1\ 9\ 10$ and $\lambda(\pi) = 1\ 1\ 2\ 4\ 4\ 2\ 1\ 1\ 9\ 1\ 9\ 10$.

Theorem 4. *The map λ from D_n to T_n^\bullet is a bijection such that*

$$\mathbf{pex}\ \pi = \mathbf{fix}\ \lambda(\pi).$$

Proof. Since the cardinality of T_n^\bullet equals that of D_n , and the image of D_n by λ is contained in T_n^\bullet , it suffices to prove the injectivity.

Let π and σ , $\pi \neq \sigma$, two derangements in D_n . If π and σ do not have the same pure excedances, then, by construction, $\lambda(\pi)$ and $\lambda(\sigma)$ do not have the same fixed points, and thus $\lambda(\pi) \neq \lambda(\sigma)$.

Now, let us assume that π and σ have the same pure excedances. If there is a pure excedance i such that $\pi^{-1}(i) \neq \sigma^{-1}(i)$ then the definition implies $\lambda(\pi) \neq \lambda(\sigma)$. Otherwise the two permutations have the same pure excedances i , and for each of them we have $\pi^{-1}(i) = \sigma^{-1}(i)$. Let j be the greatest integer such that $\pi_j \neq \sigma_j$ (without loss of generality, we assume $\pi_j < \sigma_j$). In this case, j is not a pure excedance for the two permutations. Thus, $\lambda(\pi)_j$ (resp. $\lambda(\sigma)_j$) is the sum of the label of (j, π_j) (resp. (j, σ_j)) with the number of pure excedances $k < j$ such that $\pi^{-1}(k) < j$ (resp. $\sigma^{-1}(k) < j$). Since we

have $\pi_j < \sigma_j$, the label of (j, π_j) is less than the label of (j, σ_j) , and the number of pure excedances $k < j$ such that $\pi^{-1}(k) < j$ is less than or equal to the number of pure excedances $k < j$ such that $\sigma^{-1}(k) < j$. Then we have $\lambda(\pi)_j < \lambda(\sigma)_j$. Then λ is an injective map, and thus a bijection. \square

Theorem 5. *There is a one-to-one correspondence ψ on the set D_n of n -length derangements such that for any $\pi \in D_n$,*

$$\mathbf{pex} \pi = \mathbf{cyc} \psi(\pi).$$

Proof. Considering Theorem 3 and Theorem 4, we define for any $\pi \in D_n$, $\psi(\pi) = \phi(\sigma)$ where σ is the permutation having $\lambda(\pi)$ as transposition array. \square

Theorem 6. *The two bivariate statistics $(\mathbf{pex}, \mathbf{fix})$ and $(\mathbf{pcyc}, \mathbf{fix})$ are equidistributed on S_n .*

Proof. Considering Theorem 5, we define the map $\bar{\psi}$ on S_n . Let π' be the permutation obtained from π by deleting all fixed points and after normalization as a permutation. Let $I = \{i_1, i_2, \dots, i_k\}$ be the set of fixed points of π . Then, we set $\pi'' = \psi(\pi')$. So, $\sigma = \bar{\psi}(\pi)$ is obtained from π'' by inserting fixed points $i \in I$ after a shift of all other entries in order to produce a permutation in S_n . By construction, we have $\mathbf{pex} \pi = \mathbf{pcyc} \sigma$ and $\mathbf{fix} \pi = \mathbf{fix} \sigma$ which completes the proof. \square

A byproduct of this theorem is

Corollary 2. *The statistics \mathbf{cyc} and $\mathbf{pex} + \mathbf{fix}$ are equidistributed on S_n .*

Also, a direct consequence of Theorem 3 and Theorem 6 is

Theorem 7. *The two statistics \mathbf{pex} and \mathbf{des}_2 are equidistributed on S_n .*

Notice that the Foata's first transformation is not a candidate for proving the equidistribution of \mathbf{pex} and \mathbf{des}_2 , while it transports \mathbf{exc} to \mathbf{des} .

Corollary 3. *For $0 \leq i \leq 2$, the patterns p_i and pex are equipopular on S_n (see [A001705](#) in [15]).*

Finally, we present two conjectures for future works.

Conjecture 1. *The two bivariate statistics $(\mathbf{des}_2, \mathbf{cyc})$ and $(\mathbf{pex}, \mathbf{cyc})$ are equidistributed on S_n .*

Conjecture 2. *The two bistatistics $(\mathbf{des}_2, \mathbf{des})$ and $(\mathbf{pex}, \mathbf{exc})$ are equidistributed on S_n .*

It is interesting to remark that $(\mathbf{des}, \mathbf{cyc})$ and $(\mathbf{exc}, \mathbf{cyc})$ are not equidistributed. Indeed, there are 3 permutations in S_3 having $\mathbf{exc} = 1$ and $\mathbf{cyc} = 2$, namely 132, 213, 321, but only 2 permutations with $\mathbf{des} = 1$ and $\mathbf{cyc} = 2$, videlicet 132 and 213. So, if the conjectures 1 and 2 are true then their proofs are probably independent.

Acknowledgements

We would like to greatly thank Vincent Vajnovszki for having offered us Conjecture 2.

References

- [1] J.-L. Baril. Statistics-preserving bijections between classical and cycle permutations. *Information Processing Letters*, 113, 17-22, 2013.
- [2] J.-L. Baril and S. Kirgizov. The pure descent statistic on permutations. *Discrete Mathematics*, 340(10), 2250-2558, 2017.
- [3] J.-L. Baril and V. Vajnovszki. Popularity of patterns over d -equivalence classes of words and permutations. *Theoretical Computer Science*, 814, 249-258, 2020.
- [4] F. Bergeron, N. Bergeron, R.B. Howlett and D.E. Taylor. A decomposition of the descent algebra of a finite Coxeter group. *J. Algebraic Combin.*, 1, 23-44; 1992.
- [5] M. Bóna. Surprising symmetries in objects counted by Catalan numbers. *The Electronic J. of Comb.*, 19(1):P62, 2012. (electronic).
- [6] P. Brändén and A. Claesson. Mesh patterns and the expansion of permutation statistics as sums of permutation patterns. *The Electronic J. of Combinatorics*, 18(2), P5, 2011.
- [7] S. Elizalde. Descent sets of cyclic permutations. *Adv. in Appl. Math.*, 47.4, 688–709, 2011.
- [8] S. Elizalde and J.M. Troyka. The number of cycles with a given descent set. *Séminaire Lotharingien de Combinatoire*, 80, #8, 2018.

- [9] D. Foata and M.P. Schützenberger. Théorie Géométrique des Polynômes Euleriens. *Lecture Notes in Math.*, 138, Springer-Verlag, Berlin, 1970.
- [10] D. Foata and G.-N. Han. Decreases and descents in words. *Séminaire Lotharinguien de Combinatoire*, 58, B58a, 2007.
- [11] A. Garsia and C. Reutenauer. A decomposition of Solomon’s descent algebra. *Adv. Math.*, 77, 189-262, 1989.
- [12] I. Gessel and G. Viennot. Binomial determinants, paths, and hook length formulae. *Adv. Math.*, 58:300–321, 1985.
- [13] S. Kitaev and P.B. Zhang. Distributions of mesh patterns of short lengths. *Advances in Applied Math.*, 110, 1-32, 2019.
- [14] P.A. MacMahon. *Combinatory Analysis*, Vols. 1 and 2, Cambridge Univ. Press, Cambridge, UK, 1915 (reprinted by Chelsea, New York, 1955).
- [15] N.J.A. Sloane. The On-line Encyclopedia of Integer Sequences, available electronically at <http://oeis.org>.
- [16] R.P. Stanley. *Enumerating Combinatorics*, Vol. 2, Cambridge University Press, 1999.
- [17] R.M. Wilson and J.H. van Lint. *A course in combinatorics*, Vol. I, Cambridge University Press, 2002.